

SCIENTIFIC OPINION

Scientific Opinion on Dietary Reference Values for vitamin A¹

EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA)^{2,3}

European Food Safety Authority (EFSA), Parma, Italy

ABSTRACT

Following a request from the European Commission, the Panel on Dietetic Products, Nutrition and Allergies derived Dietary Reference Values for vitamin A. The Panel considered that a concentration of 20 µg retinol/g liver can be used as a target for establishing the Average Requirement (AR) for vitamin A. In the absence of a better characterisation of the relationship between vitamin A intake and liver stores, a factorial approach was applied. This approach considered a total body/liver retinol store ratio of 1.25, a liver/body weight ratio of 2.4 %, a fractional catabolic rate of body retinol of 0.7 % per day, an efficiency of storage in the whole body for ingested retinol of 50 % and reference weights for women and men in the EU of 58.5 and 68.1 kg, respectively. ARs of 570 µg retinol equivalent (RE)/day for men and 490 µg RE/day for women were derived. Assuming a coefficient of variation (CV) of 15 %, Population Reference Intakes (PRIs) of 750 µg RE/day for men and 650 µg RE/day for women were set. For infants aged 7–11 months and children, the same equation as for adults was applied by using specific values for reference weight and liver/body weight ratio. For catabolic rate, the adult value corrected on the basis of a growth factor was used. ARs range from 190 µg RE/day in infants aged 7–11 months to 580 µg RE/day in boys aged 15–17 years. PRIs for infants and children were estimated using a CV of 15 % and range from 250 to 750 µg RE/day. For pregnancy and lactation, additional vitamin A requirements related to the accumulation of retinol in fetal and maternal tissues and transfer of retinol into breast milk were considered and PRIs of 700 and 1 300 µg RE/day, respectively, were set.

© European Food Safety Authority, 2015

KEY WORDS

vitamin A, retinol, carotenoids, Average Requirement, Population Reference Intake, Dietary Reference Value

¹ On request from the European Commission, Question No EFSA-Q-2011-01226, adopted on 5 February 2015.

² Panel members: Carlo Agostoni, Roberto Berni Canani, Susan Fairweather-Tait, Marina Heinonen, Hannu Korhonen, Sébastien La Vieille, Rosangela Marchelli, Ambroise Martin, Androniki Naska, Monika Neuhäuser-Berthold, Grażyna Nowicka, Yolanda Sanz, Alfonso Siani, Anders Sjödin, Martin Stern, Sean (J.J.) Strain, Inge Tetens, Daniel Tomé, Dominique Turck and Hans Verhagen. Correspondence: nda@efsa.europa.eu

³ Acknowledgement: The Panel wishes to thank the members of the Working Group on Dietary Reference Values for vitamins: Christel Lamberg-Allardt, Monika Neuhäuser-Berthold, Grażyna Nowicka, Kristina Pentieva, Hildegard Przyrembel, Sean (J.J.) Strain, Inge Tetens, Daniel Tomé and Dominique Turck for the preparatory work on this scientific opinion and EFSA staff: Anja Brönstrup, Agnès de Sesmaisons-Lecarré, Liisa Valsta, Fanny Héraud and Sofia Ioannidou for the support provided to this scientific opinion.

Suggested citation: EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2015. Scientific opinion on Dietary Reference Values for vitamin A. EFSA Journal 2015;13(3):4028, 84 pp. doi:10.2903/j.efsa.2015.4028

Available online: www.efsa.europa.eu/efsajournal

SUMMARY

Following a request from the European Commission, the EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) was asked to deliver a scientific opinion on Dietary Reference Values (DRVs) for the European population, including vitamin A.

Vitamin A is a fat-soluble vitamin obtained from the diet either as preformed vitamin A (mainly retinol and retinyl esters) in foods of animal origin or as provitamin A carotenoids in plant-derived foods. The term vitamin A comprises all-*trans*-retinol (also called retinol) and the family of naturally occurring molecules associated with the biological activity of retinol (such as retinal, retinoic acid, retinyl esters), as well as provitamin A carotenoids that are dietary precursors of retinol. The biological value of substances with vitamin A activity is expressed as retinol equivalent (RE). Specific carotenoids/retinol equivalency ratios are defined for provitamin A carotenoids, which account for the less efficient absorption of carotenoids and their bioconversion to retinol. On the basis of available evidence, the Panel decided to maintain the conversion factors proposed by the Scientific Committee for Food (SCF) for the European populations, namely 1 µg RE equals 1 µg of retinol, 6 µg of β-carotene and 12 µg of other provitamin A carotenoids. Vitamin A requirement can be met with any mixture of preformed vitamin A and provitamin A carotenoids that provides an amount of vitamin A equivalent to the reference value in terms of µg RE/day.

Vitamin A is involved in vision as retinal, which plays a central role in the mechanisms of photo-transduction, and in the systemic maintenance of the growth and integrity of cells in body tissues through the action of retinoic acid, which acts as regulator of genomic expression. The most specific clinical consequence of vitamin A deficiency is xerophthalmia, which encompasses a clinical spectrum of ocular manifestations. In low-income countries, vitamin A deficiency in young infants and children has been associated with increased infectious morbidity and mortality, including respiratory infection and diarrhoea.

Preformed vitamin A is efficiently absorbed (70–90 %). The absorption of β-carotene appears to be highly variable (5–65 %), depending on food- and diet-related factors, genetic characteristics and the health status of the subject. The intestine is the primary tissue where dietary provitamin A carotenoids are converted to retinol. Retinol, in the form of retinyl esters, and provitamin A carotenoids enter the body as a component of nascent chylomicrons secreted into the lymphatic system. Most dietary retinol (in chylomicrons and chylomicron remnants) is taken up by the liver, which is the major site of retinol metabolism and storage. Hepatic retinyl esters are hydrolysed to free retinol, and delivered to tissues by retinol-binding protein. The efficiency of storage and catabolism of retinol depends on vitamin A status. Low retinol stores are associated with a reduced efficiency of storage and decreased absolute catabolic rate. The majority of retinol metabolites are excreted in urine, in faeces via bile and to a lesser extent in breath.

Vitamin A status is best expressed in terms of total body store of retinol (i.e. as free retinol and retinyl esters) or, alternatively, as liver concentration of the vitamin. A concentration of 20 µg retinol/g liver (0.07 µmol/g) in adults represents a level assumed to maintain adequate plasma retinol concentration, to prevent clinical signs of deficiency and to provide adequate stores. The Panel considered that this can be used as a target value for establishing the Average Requirement (AR) for vitamin A for all age groups. The relationship between dietary intake of vitamin A and retinol liver stores has been explored with stable isotope dilution methods but available data are considered insufficient to derive an AR. A factorial approach was applied. This approach considered a total body/liver retinol store ratio of 1.25 (i.e. 80 % of retinol body stores are in the liver), a liver/body weight ratio of 2.4 %, a fractional catabolic rate of retinol of 0.7 % per day of total body stores, an efficiency of storage in the whole body of ingested retinol of 50 % and the reference body weights for women and men in the EU of 58.5 and 68.1 kg, respectively. On the basis of this approach, ARs of 570 µg RE/day for men and 490 µg RE/day for women were derived after rounding. Assuming a coefficient of variation (CV) of 15 % because of the variability in requirement and the large uncertainties in the dataset, Population

Reference Intakes (PRIs) of 750 µg RE/day for men and 650 µg RE/day for women were set after rounding.

For infants aged 7–11 months and children, the same target concentration of retinol in the liver and the same equation as for adults was used to calculate ARs. Specific values for reference body weight and for liver/body weight ratio were used. There are some indications that retinol catabolic rate may be higher in children than in adults, but data are limited. The Panel decided to apply the value for catabolic rate in adults and correct it on the basis of a growth factor. Estimated ARs range from 190 µg RE/day in infants aged 7–11 months to 580 µg RE/day in boys aged 15–17 years. PRIs for infants and children were estimated based on a CV of 15 % and range from 250 to 750 µg RE/day.

For pregnant women, the Panel assumed that a total amount of 3 600 µg retinol is accumulated in the fetus over the course of pregnancy. Considering that the accretion mostly occurs in the last months of pregnancy, and assuming an efficiency of storage of 50 % for the fetus, an additional daily requirement of 51 µg RE was calculated for the second half of pregnancy. In order to allow for the extra need related to the growth of maternal tissues, the Panel applied this additional requirement to the whole period of pregnancy. Consequently, an AR of 540 µg RE/day was estimated for pregnant women. Considering a CV of 15 % and rounding, a PRI of 700 µg RE/day was derived for pregnant women.

For lactating women, an increase in the AR was based on the vitamin A intake required to compensate for the loss of retinol in breast milk. Based on an average amount of retinol secreted in breast milk of 424 µg/day and an absorption efficiency of retinol of 80 %, an additional vitamin A intake of 530 µg RE/day was considered sufficient to replace these losses. An AR of 1 020 µg RE/day was estimated and, considering a CV of 15 % and rounding down, a PRI of 1 300 µg RE/day was proposed for lactating women.

Foods rich in retinol include offal and meat, butter, retinol-enriched margarine, dairy products and eggs, while foods rich in β-carotene include vegetables and fruits, such as sweet potatoes, carrots, pumpkins, dark green leafy vegetables, sweet red peppers, mangoes and melons. On the basis of data from 12 dietary surveys in nine EU countries, vitamin A intake was assessed using food consumption data from the EFSA Comprehensive Food Consumption Database and vitamin A composition data from the EFSA nutrient composition database. Average vitamin A intake ranged between 409 and 651 µg RE/day in children aged 1 to < 3 years, between 607 and 889 µg RE/day in children aged 3 to < 10 years, between 597 and 1 078 µg RE/day in children aged 10 to < 18 years and between 816 and 1 498 µg RE/day in adults.

TABLE OF CONTENTS

Abstract.....	1
Summary.....	2
Background as provided by the European Commission.....	6
Terms of reference as provided by the European Commission.....	6
Assessment.....	8
1. Introduction.....	8
2. Definition/category.....	8
2.1. Chemistry.....	8
2.2. Function of vitamin A.....	9
2.2.1. Biochemical functions.....	9
2.2.2. Health consequences of deficiency and excess.....	10
2.3. Physiology and metabolism.....	12
2.3.1. Intestinal absorption.....	12
2.3.2. Transport in blood.....	14
2.3.3. Distribution to tissues.....	15
2.3.4. Storage.....	15
2.3.5. Metabolism.....	17
2.3.6. Elimination.....	18
2.3.7. Interaction with other nutrients.....	20
2.3.8. Retinol equivalents.....	20
2.4. Biomarkers.....	22
2.4.1. Total body retinol content and liver retinol concentration.....	22
2.4.2. Plasma/serum retinol concentration.....	24
2.4.3. Markers of visual function.....	25
2.4.4. Conclusion on biomarkers.....	26
2.5. Effects of genotypes.....	26
3. Dietary sources and intake data.....	26
3.1. Dietary sources.....	26
3.2. Dietary intake.....	27
4. Overview of Dietary Reference Values and recommendations.....	28
4.1. Adults.....	28
4.2. Infants and children.....	30
4.3. Pregnancy.....	31
4.4. Lactation.....	32
5. Criteria (endpoints) on which to base Dietary Reference Values.....	33
5.1. Indicators of vitamin A requirement.....	33
5.1.1. Symptoms of vitamin A deficiency.....	33
5.1.2. Plasma/serum retinol concentration.....	33
5.1.3. Maintenance of total body retinol content and liver retinol concentration.....	34
5.2. Indicators of vitamin A requirement in pregnancy and lactation.....	36
5.3. Vitamin A intake and health consequences.....	37
6. Data on which to base Dietary Reference Values.....	38
6.1. Adults.....	38
6.2. Infants and children.....	39
6.3. Pregnancy.....	40
6.4. Lactation.....	40
Conclusions.....	40
Recommendations for research.....	41
References.....	42
Appendices.....	65
Appendix A. Prospective cohort and nested case–control studies on the association between intake of “vitamin A” and retinol and risk of bone fracture.....	65

Appendix B.	Intervention and prospective cohort studies on the association between intake of “vitamin A” and retinol and measures of bone mineral content (BMC), bone mineral density (BMD) or serum markers of bone turnover.....	70
Appendix C.	Retinol concentration in breast milk from mothers of term infants	73
Appendix D.	Dietary surveys in the updated EFSA Comprehensive European Food Consumption Database included in the nutrient intake calculation and number of subjects in the different age classes.....	75
Appendix E.	Vitamin A intake in males in different surveys according to age classes and country ($\mu\text{g RE/day}$).....	76
Appendix F.	Vitamin A intake in females in different surveys according to age classes and country ($\mu\text{g RE/day}$).....	78
Appendix G.	Minimum and maximum % contribution of different food groups to vitamin A intake in males	80
Appendix H.	Minimum and maximum % contribution of different food groups to vitamin A intake in females	81
Abbreviations.....		82

BACKGROUND AS PROVIDED BY THE EUROPEAN COMMISSION

The scientific advice on nutrient intakes is important as the basis of Community action in the field of nutrition, for example such advice has in the past been used as the basis of nutrition labelling. The Scientific Committee for Food (SCF) report on nutrient and energy intakes for the European Community dates from 1993. There is a need to review and, if necessary, to update these earlier recommendations to ensure that the Community action in the area of nutrition is underpinned by the latest scientific advice.

In 1993, the SCF adopted an opinion on the nutrient and energy intakes for the European Community.⁴ The report provided Reference Intakes for energy, certain macronutrients and micronutrients, but it did not include certain substances of physiological importance, for example dietary fibre.

Since then new scientific data have become available for some of the nutrients, and scientific advisory bodies in many European Union Member States and in the United States have reported on recommended dietary intakes. For a number of nutrients these newly established (national) recommendations differ from the reference intakes in the SCF (1993) report. Although there is considerable consensus between these newly derived (national) recommendations, differing opinions remain on some of the recommendations. Therefore, there is a need to review the existing EU Reference Intakes in the light of new scientific evidence, and taking into account the more recently reported national recommendations. There is also a need to include dietary components that were not covered in the SCF opinion of 1993, such as dietary fibre, and to consider whether it might be appropriate to establish reference intakes for other (essential) substances with a physiological effect.

In this context, EFSA is requested to consider the existing Population Reference Intakes for energy, micro- and macronutrients and certain other dietary components, to review and complete the SCF recommendations, in the light of new evidence, and in addition advise on a Population Reference Intake for dietary fibre.

For communication of nutrition and healthy eating messages to the public it is generally more appropriate to express recommendations for the intake of individual nutrients or substances in food-based terms. In this context, EFSA is asked to provide assistance on the translation of nutrient based recommendations for a healthy diet into food based recommendations intended for the population as a whole.

TERMS OF REFERENCE AS PROVIDED BY THE EUROPEAN COMMISSION

In accordance with Article 29(1)(a) and Article 31 of Regulation (EC) No 178/2002,⁵ the Commission requests EFSA to review the existing advice of the Scientific Committee for Food on population reference intakes for energy, nutrients and other substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

In the first instance, EFSA is asked to provide advice on energy, macronutrients and dietary fibre. Specifically advice is requested on the following dietary components:

- Carbohydrates, including sugars;
- Fats, including saturated fatty acids, polyunsaturated fatty acids and monounsaturated fatty acids, *trans* fatty acids;

⁴ Scientific Committee for Food, Nutrient and energy intakes for the European Community, Reports of the Scientific Committee for Food 31st series, Office for Official Publication of the European Communities, Luxembourg, 1993.

⁵ Regulation (EC) No 178/2002 of the European Parliament and of the Council of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety. OJ L 31, 1.2.2002, p. 1–24.

- Protein;
- Dietary fibre.

Following on from the first part of the task, EFSA is asked to advise on population reference intakes of micronutrients in the diet and, if considered appropriate, other essential substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

Finally, EFSA is asked to provide guidance on the translation of nutrient based dietary advice into guidance, intended for the European population as a whole, on the contribution of different foods or categories of foods to an overall diet that would help to maintain good health through optimal nutrition (food-based dietary guidelines).

ASSESSMENT

1. Introduction

In 1993, the Scientific Committee for Food (SCF) adopted an opinion on nutrient and energy intakes for the European Community and derived Average Requirements (ARs) and Population Reference Intakes (PRIs) for vitamin A for men and women. Specific PRIs were set for pregnant and lactating women. PRIs for infants aged 7–11 months and children were also proposed. Vitamin A is a fat-soluble vitamin obtained from the diet either as preformed vitamin A (mainly retinol and retinyl esters) in foods of animal origin, or as provitamin A carotenoids in plant-derived foods (Figure 1). The purpose of this opinion is to review Dietary Reference Values (DRVs) for vitamin A. The Panel notes that possible functions of carotenoids other than as dietary precursors of retinol, and evidence for a requirement for carotenoids as such, have been reviewed by the SCF (1993) and other authoritative bodies (DH, 1991; IOM, 2001; WHO/FAO, 2004; D-A-CH, 2013). This is out of the scope of the present opinion.

2. Definition/category

2.1. Chemistry

The term vitamin A comprises all-*trans*-retinol (also called retinol) and the family of naturally occurring molecules associated with the biological activity of retinol (such as retinal, retinoic acid and retinyl esters), as well as the group of provitamin A carotenoids (such as β -carotene, α -carotene and β -cryptoxanthin) that are dietary precursors of retinol (Figure 1).




Figure 1: Structure of the naturally occurring forms of vitamin A: all-*trans*-retinol, an all-*trans*-retinyl ester (R=alkyl chain), all-*trans*-retinal, the active metabolites all-*trans*-retinoic acid (transcriptionally active) and 11-*cis*-retinal (active in vision), and the major provitamin A carotenoid, all-*trans*- β -carotene.

Retinol is composed of a β -ionone ring, a polyunsaturated side chain and a polar end group (molecular mass 286.5 Da) (Figure 1). This chemical structure makes it poorly soluble in water but easily transferable through membrane lipid bilayers. Preformed vitamin A consists predominantly of retinol and retinyl esters, which are supplied in the diet by animal-derived products. The term retinoids refers

to retinol and structurally related compounds, including its metabolites (retinyl ester, retinal and retinoic acid), and synthetic analogues (Anonymous, 1983).

Retinol and retinyl esters are the most abundant forms of vitamin A in the body. Retinol is a transport form and a precursor of the transcriptionally active metabolite all-*trans*-retinoic acid, and retinyl esters are retinol storage forms and serve as substrate for the formation of the visual chromophore 11-*cis*-retinal (Al Tanoury et al., 2013). All-*trans*-retinoic acid can be isomerised through a non-enzymatic process to 9-*cis*- or 13-*cis*-retinoic acid isomers. The isomer 13-*cis*-retinoic acid is less transcriptionally active than the all-*trans* and the 9-*cis* isomers. Other forms of retinol and retinoic acid, which include various oxo-, hydroxy- and glucuronide forms, are also present in the body, but at very low concentrations relative to retinol and retinyl esters, and probably appear as catabolic products for elimination from the body (O'Byrne and Blaner, 2013).

Carotenoids are isoprenoids that contain up to 15 conjugated double bonds, synthesised in plants and microorganisms and occurring naturally in fruits and vegetables. Among them, β -carotene, α -carotene, and β -cryptoxanthin are provitamin A carotenoids (Eroglu and Harrison, 2013). To exhibit provitamin A activity, the carotenoid molecule must have at least one unsubstituted β -ionone ring and the correct number and position of methyl groups in the polyene chain (Wirtz et al., 2001).

In this opinion, the terms retinol, retinoic acid and carotenoids refer to their all-*trans*-isomers, unless specified otherwise.

The biological value of substances with vitamin A activity is expressed as retinol equivalent (RE), with 1 μg RE equal to 1 μg retinol. Specific carotenoids/retinol equivalency ratios are defined for provitamin A carotenoids, which account for the less efficient absorption of carotenoids and their bioconversion to retinol (Section 2.3.8).

2.2. Function of vitamin A

2.2.1. Biochemical functions

Vitamin A is an essential nutrient as humans do not have the capability for *de novo* synthesis of compounds with vitamin A activity. Vitamin A is involved in the visual cycle in the retina and the systemic maintenance of growth and integrity of cells in body tissues.

In the eye, the active metabolite 11-*cis*-retinal works as a visual chromophore involved in phototransduction. Visual pigments are G-protein-coupled receptors that mediate phototransduction, the process by which light is translated into an electrical (nervous) signal (Palczewski, 2010). In this complex pathway, also known as the retinoid cycle, 11-*cis*-retinal binds opsin to form rhodopsin and cone pigments (Wald, 1968). Visual perception starts with the absorption of a photon, which induces isomerisation of 11-*cis*-retinal to 11-*trans*-retinal. After bleaching, 11-*trans*-retinal is released from opsin and the 11-*cis*-retinal chromophore is regenerated to sustain vision (von Lintig et al., 2010). In addition, all-*trans*-retinoic acid is also required to maintain normal differentiation of the cornea and conjunctival membranes and of the photoreceptor rod and cone cells of the retina (Blomhoff and Blomhoff, 2006).

Retinoic acid is a transcriptionally active metabolite and is thought to account for the regulatory properties of vitamin A upon more than 500 different target genes involved in the differentiation and development of fetal and adult tissues, stem cell differentiation, apoptosis, support of reproductive and immune functions and regulation of lipid metabolism and energy homeostasis (Al Tanoury et al., 2013; Kedishvili, 2013). Retinoic acid can activate two different types of nuclear receptors, retinoic acid receptor (RAR) and peroxisome proliferator-activated receptor (PPAR) β/δ . In the cytosol, retinoic acid binds to cellular retinoic acid-binding protein (CRABP) II, and the resulting complex channels retinoic acid to RARs. RARs work as heterodimers with retinoic X receptors (RXRs) and transduce the retinoic acid signal as ligand-dependent regulators of transcription. Retinoic acid also binds to fatty acid-binding protein (FABP) 5 and activates the nuclear translocation of FABP5, which

then delivers the ligand to the PPAR β/δ subtype. In addition, retinoic acid has extranuclear, non-transcriptional effects, such as the activation of the mitogen-activated protein kinase signalling pathway, which influences the expression of retinoic acid target genes via phosphorylation processes (Al Tanoury et al., 2013).

The importance of vitamin A in immune function is well established (Stephensen, 2001; Field et al., 2002). Mechanisms by which vitamin A may modulate the immune system have been studied *in vitro* and in animal models. Retinoic acid stimulates the proliferation of T-lymphoid cells, inhibits the proliferation of B-cells and B-cell precursors and exerts an effect on the T-helper (Th) cell balance by suppressing Th1 development and enhancing Th2 development. It also enhances macrophage-mediated inflammation by increasing production of interleukin 12 (IL-12) and interferon-gamma (IFN- γ), regulates the antigen presentation by immature dendritic cells, as well as their maturation, and impairs the ability of macrophages to ingest and kill bacteria (Ross et al., 2011; Cassani et al., 2012; Ross, 2012). Other effects of vitamin A on the immune system are related to apoptotic effects on immune-competent cells during down-regulation of immune reactions and during thymic selection and to the alteration of genes relevant to the immune response (Ruhl, 2007).

2.2.2. Health consequences of deficiency and excess

2.2.2.1. Deficiency

The main symptoms observed in case of deficiency of vitamin A are intrauterine and post-natal growth retardation and a large array of congenital malformations collectively referred to as the fetal “vitamin A deficiency syndrome”, which is well documented in animals (Clagett-Dame and Knutson, 2011). In adults, vitamin A deficiency adversely affects several functions, such as vision, immunity, and reproduction, and has been related to the worsening of low iron status, resulting in vitamin A deficiency anaemia (Ross, 2014).

The most specific clinical consequence of vitamin A deficiency is xerophthalmia, which encompasses the clinical spectrum of ocular manifestations of vitamin A deficiency. It includes night blindness (nyctalopia), due to impaired dark adaptation because of slow regeneration of rhodopsin, early keratinising metaplasia (Bitot’s spots), impaired production of tears, conjunctival xerosis, corneal xerosis, and corneal ulceration and scarring, which may result in blindness (WHO, 1982, 1996, 2009). Night blindness, the first ocular symptom of deficiency, responds rapidly to an increase in vitamin A intake (Dowling and Gibbons, 1961; Sommer A, 1982; Katz et al., 1995; Christian et al., 1998a).

Vitamin A deficiency also induces follicular hyperkeratosis, which disappears after retinol or β -carotene supplementation (Chase et al., 1971; Sauberlich et al., 1974).

In low-income countries, vitamin A deficiency in young infants and children has been associated with increased infectious morbidity and mortality, including respiratory infection and diarrhoea (Mayo-Wilson et al., 2011).

2.2.2.2. Excess

The classical signs and symptoms of acute and chronic hypervitaminosis A comprise skin disorders, nausea, vomiting, disorders of the musculo-skeletal system and liver damage (Biesalski, 1989; Hathcock et al., 1990). Bulging fontanelle in infants and increased intracranial pressure are also classical adverse effects of vitamin A toxicity (Hathcock et al., 1990). The teratogenic effect of excessive intake of vitamin A or specific retinoids is well documented, in both animals and humans (Hathcock et al., 1990).

In 2002, the SCF reviewed possible adverse effects of long-term intake of retinol and retinyl esters (SCF, 2002). The SCF set a Tolerable Upper Intake Level (UL) for preformed vitamin A of 3 000 $\mu\text{g RE/day}$ for women of childbearing age and men, based on the risk of hepatotoxicity and teratogenicity. The UL was proposed to also apply during pregnancy and lactation. ULs for children

were extrapolated from the UL for adults, based on allometric scaling (body weight to the power of 0.75). ULs were set at 800 µg RE/day for children aged 1–3 years, 1 100 µg RE/day for children aged 4–6 years, 1 500 µg RE/day for children aged 7–10 years, 2 000 µg RE/day for children aged 11–14 years and 2 600 µg RE/day for children aged 15–17 years.

The SCF noted that an increased risk of bone fracture was reported for an intake of 1 500 µg RE/day or higher. Presumed mechanisms related to a possible effect of retinoic acid on osteoblasts and osteoclasts (Scheven and Hamilton, 1990; Kindmark et al., 1995; Cohen-Tanugi and Forest, 1998) and a molecular interaction of vitamin A and vitamin D indicating an antagonism of vitamin A towards the action of vitamin D (Rohde et al., 1999; Johansson and Melhus, 2001) were mentioned. Overall, the SCF considered that the available data did not provide sufficient evidence of causality, owing to the possibility of residual confounding, and were not appropriate for establishing a UL. The SCF noted that *“because the tolerable upper level may not adequately address the possible risk of bone fracture in particularly vulnerable groups, it would be advisable for postmenopausal women, who are at greater risk of osteoporosis, to restrict their intake to 1 500 µg RE/day”*.

In a subsequent assessment which considered studies published until 2004, the Scientific Advisory Committee on Nutrition (SACN, 2005) concluded that the evidence of an association between high intake of retinol and poor bone health was inconsistent. The Committee noted that some epidemiological data suggest that a retinol intake of 1 500 µg/day and above is associated with an increased risk of bone fracture; the evidence was considered not robust enough to set a Safe Upper Level, and a Guidance Level for retinol intake of 1 500 µg/day was set for adults.

The Panel is aware that additional observational studies on possible associations between retinol and vitamin A intake and bone health have been published since the SCF and SACN assessments. The Panel notes that different definitions of “vitamin A” have been applied among studies (i.e. defined as retinol only, as retinol and provitamin A carotenoids (expressed in IU⁶ or µg RE) or undefined). An overview of prospective cohort and nested case–control studies which investigated an association of retinol or “vitamin A” intake with the risk of bone fracture is provided in Appendix A. Intervention and prospective cohort studies which looked at an association of retinol or “vitamin A” intake with markers of bone health are summarised in Appendix B. These appendices tabulate studies considered in the SCF and SACN assessments, along with studies published afterwards. Among the latter, no association was observed between a cumulative dose of retinol supplementation and the risk of any fracture or “osteoporotic fracture” (defined as fractures at the spine, hip, femur, arm, ribs or wrist) in 2 322 Australian males and females who received 7.5 mg RE/day as retinyl palmitate for 1 to 16 years (187 subjects experienced 237 fractures) (Ambrosini et al., 2013). Similarly, no association was found between retinol or “vitamin A” intake (from food and supplements) and risk of any fracture or hip fracture in the Women’s Health Initiative prospective study, which involved 75 747 postmenopausal women in the USA (mean follow-up 6.6 years; 10 405 incident total fractures and 588 hip fractures). In contrast, in a stratified analysis, modest increases in total fracture risk with high retinol intake (fifth quintile (Q5) = 2 488 µg/day vs. first quintile (Q1) = 348 µg/day) (hazard ratio (HR) = 1.15; 95 % confidence interval (CI) 1.03–1.29; p for trend = 0.056) and high “vitamin A” intake (Q5 = 8 902 µg RE/day vs. Q1 = 4 445 µg RE/day) (HR = 1.19; 95 % CI 1.04–1.37; p for trend = 0.022) were observed in the women with a vitamin D intake ≤ 11 µg/day (Caire-Juvera et al., 2009). No association between retinol or “vitamin A” intake (from food only or food and supplements) and fracture risk was found in a nested case–control analysis of the Danish Osteoporosis Prevention Study which involved 1 141 perimenopausal women (163 cases, 978 controls) (Rejnmark et al., 2004). Two studies investigated bone mineral density (BMD) as an endpoint. No significant association was observed between BMD change and retinol or “vitamin A” intake (from food and supplements) in 891 women followed for five to seven years in the Aberdeen Prospective Osteoporosis Study (Macdonald et al., 2004). No association between retinol or “vitamin A” intake (from food only or food and

⁶ 1 IU = 0.3 µg retinol = 0.6 µg β-carotene = 1.2 µg other total mixed carotenoids with vitamin A activity (US DH/FAO, 1968).

supplements) and BMD or change in BMD after a five-year follow-up was found in the Danish Osteoporosis Prevention Study, which involved 1 694 women (Rejnmark et al., 2004).

The Panel is aware of other studies which investigated the association between serum/plasma retinol concentration and fracture risk (Opotowsky and Bilezikian, 2004; Ambrosini et al., 2014). Although serum/plasma retinol concentration has been used as a biomarker of intake, serum/plasma retinol concentration is under homeostatic control and, in the usual range, is not related to observed levels of habitual vitamin A intake. Therefore, it is not considered a reliable marker of vitamin A or retinol intake (Section 2.4.2).

The Panel considers that evaluation of the data published since the SCF assessment does not change the conclusion of the Panel from that of the SCF with respect to the association between retinol or vitamin A intake and risk of bone fracture in postmenopausal women. One prospective cohort study indicated a possible interaction between vitamin D intake (< 11 µg/day) and retinol intake in relation to the risk of bone fracture in postmenopausal women.

The Panel is aware of other studies which looked at possible associations between preformed vitamin A intake or blood retinol concentration and adverse health outcomes (Grotto et al., 2003; Bjelakovic et al., 2008; Chen et al., 2008; Mayo-Wilson et al., 2011; Beydoun et al., 2012; Bjelakovic et al., 2012; Bjelakovic et al., 2013; Field et al., 2013; Bjelakovic et al., 2014). Available data on individual outcomes are limited or relate to interventions that used large doses of retinol (≥ 6 000 µg) once or several times a year, which are difficult to relate to a potential effect of daily dietary intake of retinol.

2.3. Physiology and metabolism

The different forms of vitamin A undergo a complex metabolic fate with an exchange between the intestine, the plasma, the liver and other peripheral tissues (Figure 2).




Figure 2: Vitamin A forms and metabolic fates.

2.3.1. Intestinal absorption

The key digestive processes that occur within the lumen of the intestine include the release of preformed vitamin A and provitamin A carotenoids from the food matrix and their emulsification with dietary fatty acids and bile acids (Parker, 1996). The presence of dietary fat in the intestine usually

increases intestinal absorption of preformed vitamin A and provitamin A carotenoids by enhancing the secretion of pancreatic enzymes and of bile salts that provide components (lysophospholipids, monoglycerides, free fatty acids) to form luminal mixed micelles of lipids and for intracellular assembly of chylomicrons (Roels et al., 1958; Roels et al., 1963; Reddy and Srikantia, 1966; Figueira et al., 1969; Jayarajan et al., 1980; Borel et al., 1997; Jalal et al., 1998; Li and Tso, 2003; Unlu et al., 2005).

2.3.1.1. Absorption of preformed vitamin A

Preformed vitamin A is efficiently absorbed in the intestine, in the range of 70–90 % (Reddy and Sivakumar, 1972; Sivakumar and Reddy, 1972; Kusin et al., 1974). Almost complete absorption was observed in five healthy Indian children administered 1 000 µg retinyl acetate in oil (Sivakumar and Reddy, 1972). Absorption remains high even if the amount of ingested preformed vitamin A increases (Olson, 1972). A fractional absorption of around 70 % was observed in Indian children when a single high dose of retinyl acetate (60 000 µg) was administered (Reddy and Sivakumar, 1972; Kusin et al., 1974). Quantitative data on the absorption of preformed vitamin A from the diet are scarce.

Dietary retinyl esters are unable to enter the intestinal mucosa and must first be hydrolysed by retinyl ester hydrolases to yield free retinol (Harrison, 2012). Retinyl esters can be hydrolysed within the intestinal lumen by non-specific pancreatic enzymes, such as pancreatic triglyceride lipase and cholesteryl ester hydrolase, or at the mucosal cell surface by a brush border retinyl ester hydrolase (Erlanson and Borgstrom, 1968; Rigtrup and Ong, 1992; Rigtrup et al., 1994; van Bennekum et al., 2000; Reboul et al., 2006).

Free retinol is taken up into the intestinal cells by protein-mediated facilitated diffusion and passive diffusion mechanisms via the action of membrane-bound lipid transporters involved in fatty acid and cholesterol uptake. These include scavenger receptor class B, type 1 (SR-B1), CD36, NPC1L1 and a variety of ABC transporters (Hollander and Muralidhara, 1977; Hollander, 1981; Glatz et al., 1997; Abumrad et al., 1998; van Heek et al., 2001; Turley and Dietschy, 2003; Wang, 2003; Altmann et al., 2004; Davis et al., 2004; Nieland et al., 2004; During et al., 2005; Iqbal and Hussain, 2009). Free retinol then binds to specific cytoplasmic retinol-binding proteins (RBPs), i.e. the cellular retinol-binding proteins CRBPI and CRBP II (Ong, 1994). CRBP II is present at high concentrations in the enterocytes and appears to be uniquely suited for retinol absorption by the intestine (Herr and Ong, 1992; Ong, 1994; Li and Norris, 1996; Newcomer et al., 1998).

CRBP-bound retinol undergoes esterification with long-chain fatty acids, particularly with palmitic acid, catalysed mainly (about 90 %) by lecithin:retinol acyltransferase (LRAT), and to a lesser extent by the intestinal acyl-CoA:retinol acyltransferase (DGAT1) (Huang and Goodman, 1965; MacDonald and Ong, 1988; O'Byrne et al., 2005; Harrison, 2012). The resulting retinyl esters are then packed, along with dietary fat and cholesterol, into nascent chylomicrons, which are secreted into the lymphatic system for delivery to the blood (Olson, 1989; Blomhoff et al., 1991; Parker, 1996; Harrison, 2012).

2.3.1.2. Absorption of provitamin A carotenoids

Dietary provitamin A carotenoids are absorbed via passive diffusion or taken up by the enterocyte through facilitated transport via SR-B1 (van Bennekum et al., 2005; During and Harrison, 2007; Moussa et al., 2008; Harrison, 2012; von Lintig, 2012).

Once inside the enterocyte, the major part (more than 60 %) of the absorbed provitamin A carotenoids are cleaved at their central double bond (15,15') by β,β -carotene-15,15'-monooxygenase 1 (BCMO1) into all-*trans*-retinal (Devery and Milborrow, 1994; Nagao et al., 1996; Lindqvist and Andersson, 2002). All-*trans*-retinal binds to CRBP II, is incorporated intact with dietary fat and cholesterol into nascent chylomicrons, or is further oxidised irreversibly to retinoic acid or reduced reversibly to retinol (Harrison, 2012).

Less than 40 % of absorbed provitamin A carotenoids are not cleaved in the intestine (Castenmiller and West, 1998) and are absorbed intact. Along with other lipid-soluble compounds, they are incorporated into chylomicrons for transport to the liver and other tissues and are found associated with circulating lipoproteins (Johnson and Russell, 1992). The Panel notes that, because of physiological differences in provitamin A carotenoid absorption between rodents and humans, rodents are not good animal models for studying human carotenoid absorption (Huang and Goodman, 1965).

Overall, the absorption of β -carotene appears to be highly variable (5–65 %), depending on food- and diet-related factors, genetic characteristics and the health status of the subject (Haskell, 2012). This has significant implications on the bioequivalence of β -carotene to retinol (Section 2.3.8). Quantitative data on absorption of the other provitamin A carotenoids, α -carotene and β -cryptoxanthin, are more limited.

2.3.2. Transport in blood

A number of different forms of vitamin A are found in the circulation, and these differ in the fasting and postprandial states (O'Byrne and Blaner, 2013). They include retinyl esters in chylomicrons, chylomicron remnants, very low-density lipoprotein (VLDL), low-density lipoprotein (LDL) and high-density lipoprotein (HDL); retinol bound to retinol-binding protein (RBP4); retinoic acid bound to albumin; and the water-soluble β -glucuronides of retinol and retinoic acid. Provitamin A carotenoids may be absorbed intact by the intestine (Section 2.3.1) and can be found in the blood bound to chylomicrons and their remnants, VLDL, LDL and HDL (Redlich et al., 1996; Redlich et al., 1999). LDL is the major carrier of β -carotene in the plasma of fasting individuals (Romanchik et al., 1995).

Approximately two-thirds of absorbed retinol is delivered to the blood via the lymph in esterified form as retinyl palmitate and other retinyl esters present in chylomicrons. Around one-third is secreted directly into the portal circulation, probably as free retinol (Blomhoff et al., 1990; Blomhoff et al., 1991; Kane and Havel, 1995; Lemieux et al., 1998; Nayak et al., 2001).

Mean fasting concentration of retinyl esters has been reported to range from 10 to 40 $\mu\text{g/L}$ in adults (Bankson et al., 1986; Hartmann et al., 2001). In the postprandial circulation, the concentration of retinyl esters increases. Following consumption of a retinol-rich meal (~ 1 – 1.5 mg/kg body weight), mean retinyl palmitate concentration in plasma was observed to reach 7–9 $\mu\text{mol/L}$ in male volunteers (Arnhold et al., 1996; Relas et al., 2000).

In the fasting circulation, retinol bound to RBP4 is the predominant form of retinoids, with concentrations ranging from 2–4 $\mu\text{mol/L}$ in adults from Western countries (Chuwers et al., 1997; Hartmann et al., 2001). The retinol–RBP4 complex binds another plasma protein, transthyretin, which stabilises the complex and reduces renal filtration of retinol (van Bennekum et al., 2001).

Retinoic acid is present in the circulation in both the fasting and postprandial states, when it is bound to albumin. Immediately following consumption of a meal rich in retinol and retinyl esters (~ 1 mg/kg body weight), mean plasma concentration of retinoic acid increased to 254 nmol/L but was quickly restored to fasting concentrations of 14 nmol/L in 10 male volunteers (Arnhold et al., 1996).

Plasma/serum concentrations of retinyl- and retinoyl- β -glucuronides have been reported to be in the range of 5–15 nmol/L (Barua and Olson, 1986; Barua et al., 1989). Although it has been proposed that retinyl- and retinoyl- β -glucuronides, which are known to be readily hydrolysed by a number of β -glucuronidases, may serve as sources of retinoids for tissues, it is generally believed that these fully water-soluble metabolites are filtered in the kidney and eliminated quickly from the body.

Average fasting blood concentrations of β -carotene in the range 0.2–0.7 $\mu\text{mol/L}$ have been reported in adult European populations (Al-Delaimy et al., 2004; Hercberg et al., 2004). A dose–response relationship between carotenoid intake and appearance in plasma has been shown (Rock et al., 1992). Individuals consuming 30 mg/day of supplemental β -carotene for five years had a mean fasting β -carotene concentration of 3.75 $\mu\text{mol/L}$ (Redlich et al., 1999).

2.3.3. Distribution to tissues

The delivery of retinoids to tissues involves many different forms and carriers (Paik et al., 2004; O'Byrne and Blaner, 2013). Quantitatively the two most important pathways are those involving retinol bound to RBP4 and the postprandial delivery pathway.

When needed, hepatic retinyl esters are hydrolysed to free retinol, which is mobilised from the liver bound to its plasma transport protein, RBP4. Retinol–RBP4 is secreted from the liver into the circulation as a means of delivering retinol to peripheral tissues (Goodman et al., 1965; Soprano and Blaner, 1994; Quadro et al., 1999; Packer, 2005). Liver is the major site of synthesis of RBP4, but other tissues, including adipose tissue, kidney, lung, heart, skeletal muscle, spleen, eyes and testes, also express RBP4, which may be important for recycling retinoids from peripheral tissues back to the liver (Blomhoff et al., 1991). Studies on intestinal cells indicate that retinol enters by diffusion, and this is probably true for other cell types (During et al., 2002; During and Harrison, 2004; During et al., 2005; During and Harrison, 2007), although part of the uptake of RBP-bound retinol into specific target tissues is mediated by a cell surface receptor for RBP4 termed STRA6 (stimulated by retinoic acid 6). STRA6 is expressed on the surface of cells of several organs such as Sertoli cells, yolk sac and chorioallantoic placenta, choroid plexuses and retinal pigmented epithelial cells (Bouillet et al., 1997; Lewis et al., 2002; Blaner, 2007; Kawaguchi et al., 2007; Pasutto et al., 2007; Berry et al., 2013).

In the postprandial delivery pathway, retinyl esters in chylomicrons in the circulation are taken up by tissues as the chylomicron undergoes lipolysis and remodelling. Approximately 66–75 % of chylomicron retinyl esters are cleared by the liver in the rat, with the remainder cleared by peripheral tissues (Goodman et al., 1965; Blaner et al., 1994; van Bennekum et al., 1999). Postprandial unesterified retinol taken up by cells is thought to bind immediately to CRBPs that are present in tissues (Noy and Blaner, 1991). Once retinol is formed upon retinyl ester hydrolysis within the hepatocyte, it is quickly bound by apo-CRBPI, which is in molar excess of retinol in these cells (Harrison et al., 1987).

In humans, the two major sites of conversion of provitamin A carotenoids to retinoids appear to be the intestine (Section 2.3.1) and the liver (Harrison, 2012). Provitamin A carotenoids in VLDL and LDL are presumably taken up along with the lipoprotein particles by their cell surface receptors. Many tissues, including the liver, intestine, kidney, skin, skeletal muscle, adrenal gland, pancreas, testis, ovary, prostate, endometrium, mammary tissue, eyes and the mammalian embryo, express the enzyme BCMO1 and are all capable of converting β -carotene to retinoids (Yan et al., 2001; Lindqvist and Andersson, 2002; Chichili et al., 2005; Lindqvist et al., 2005). The maximum capacity for β -carotene cleavage by the intestine and the liver combined was estimated to be 12 mg β -carotene/day in a human adult (During et al., 2001). The liver and the intestine were both shown to have substantial β -carotene 15,15'-oxidoreductase activity (During et al., 2001). A study using a compartmental modelling approach showed that both the intestine and the liver should be considered to fit the prediction of the model to the experimental data for *in vivo* β -carotene conversion to retinol (Novotny et al., 1995). A study using stable isotope-labelled β -carotene (dose of 6 mg) showed that, on average, in a well-nourished population, 81 % of total vitamin A formed from a β -carotene dose is from intestinal cleavage and 19 % is from extra-intestinal cleavage (post-absorptive or whole body) (Tang et al., 2003).

Plasma retinoic acid may also be taken up into tissues through a “flip-flop” mechanism across phospholipid bilayers (Noy, 1992b, 1992a) and contribute to tissue pools (Kurlandsky et al., 1995).

2.3.4. Storage

The main storage form of retinol is retinyl esters. The liver and intestine are the major tissue sites of retinol esterification, but other tissues, including the eye, lung, adipose tissue, testes, skin and spleen, are also able to esterify retinol and accumulate retinyl ester stores. The enzyme responsible for most retinyl ester formation is LRAT. Liver LRAT is thought to be structurally identical to intestinal LRAT (Section 2.3.1.1), although hepatic but not intestinal LRAT expression appears to be regulated by the

vitamin A nutritional status (Matsuura and Ross, 1993). The concentration of retinoic acid in tissues is generally very low, usually 100 to 1 000 times less than that of retinol and retinyl esters.

2.3.4.1. Liver stores

It is considered that in healthy individuals with an adequate vitamin A status, 70–90 % of retinol of the body is stored in the liver and that this percentage decreases to 50 % or below in severely deficient individuals (Rietz et al., 1973; Bausch and Rietz, 1977; Olson, 1987). Based on empirical data, Rietz et al. (1973) indicated that, in rats with an adequate vitamin A intake, 80 % of the retinol content of the body is stored in the liver. There is a lack of direct measurement in humans. Using stable isotope and model-based compartmental analysis to study retinol kinetics in one healthy human volunteer in the USA, von Reinersdorff et al. (1998) predicted that 80 % of the absorbed dose of labelled retinol was contained in the liver seven days after administration.

The major part of retinoids is concentrated in the lipid droplets of hepatic stellate cells (Hendriks et al., 1985; Moriwaki et al., 1988; Blomhoff et al., 1991; Blaner et al., 2009), where nearly all of the retinoids present is stored as retinyl ester (primarily retinyl palmitate, with smaller amounts of retinyl stearate, retinyl oleate and retinyl linoleate) (Blaner et al., 1985; Blomhoff et al., 1991; Blaner et al., 2009). Unesterified retinol accounts for less than 1 %.

Hepatocytes are responsible for the uptake of chylomicron remnant retinoids into the liver, which are then transferred to hepatic stellate cells (Blaner et al., 1985; Blomhoff et al., 1991). Hepatocytes account for about 10–20 % of the retinoids stored in the liver (Blaner et al., 1985; Blaner et al., 2009). Hepatocytes are the sole hepatic cellular site of RBP4 synthesis and possess enzymatic activities needed for the hydrolysis of retinyl esters and the synthesis and catabolism of retinoic acid (Blaner et al., 1985; Blaner et al., 2009).

2.3.4.2. Adipose tissue stores

Adipocytes are able to accumulate significant retinyl ester stores (O'Byrne and Blaner, 2013). Data in rats indicate that the adipose tissue may account for as much as 15–20 % of the total body retinoids (Tsutsumi et al., 1992). Data in humans are lacking. As in the liver, retinyl esters stored in adipose tissue can be mobilised and secreted back into the circulation bound to RBP4 synthesised in adipocytes (Tsutsumi et al., 1992; Zovich et al., 1992; Wei et al., 1997). These retinyl esters are first hydrolysed by hormone-sensitive lipase, which acts as retinyl ester hydrolase in adipocytes (Wei et al., 1997; Strom et al., 2009).

2.3.4.3. Efficiency of storage

The efficiency of storage represents the fraction of ingested retinol which is absorbed and retained in the body (and more particularly in the liver).

Upon i.v. administration of ³H-labelled retinol to rats with different vitamin A stores, the percentage of storage in the liver was shown to be relatively constant, between 50 and 63 %, in the range of liver retinol concentrations of 18–54 µg/g (Bausch and Rietz, 1977). The percentage of ³H-labelled retinol stored in the liver decreased (6–40 %) when initial hepatic stores of retinol were below 18 µg/g liver (0.06 µmol/g) (Bausch and Rietz, 1977).

Using a radio-isotopic method, whole-body retinol retention was assessed in groups of Indian children (2–10 years) by measuring radioactivity in urine and faeces over four to six days after administration of a labelled dose (Reddy and Sivakumar, 1972; Sivakumar and Reddy, 1972). When the labelled dose of retinyl acetate was administered with 1 000 µg unlabelled retinyl acetate, mean retention was 82.2 ± 2.0 % in healthy children (n = 5) and 57.6 ± 6.0 % in a group of children (n = 8) with infection. When the labelled dose of retinyl palmitate was administered with a high dose of 60 000 µg retinyl palmitate in five healthy children, 47 % of the dose was retained, on average. Using similar methodology, retention in the range of 48–54 % was estimated in healthy Indian children (n = 17; 3–6 years), when

labelled retinyl acetate was administered with a high dose of 60 000 µg unlabelled retinyl acetate (Kusin et al., 1974). The liver retinol content of these children is unknown.

The efficiency of storage in the liver was measured by stable isotopic methods in 31 adult Bangladeshi surgery patients. Liver biopsy samples were taken 9–11 days after the administration of an oral dose of labelled retinol (215 µg/kg body weight (0.753 µmol/kg body weight) as retinyl acetate) (Haskell et al., 1997). In patients with hepatic concentrations greater than or equal to 20 µg retinol/g of liver (0.07 µmol/g) (mean ± SD estimated hepatic stores 40 ± 18 mg RE (139 ± 64 µmol RE)), average efficiency of storage in the liver was 42 % (± 13 %). The efficiency of storage in the liver was significantly lower (30 ± 8 %) in subjects with a concentration < 20 µg retinol/g of liver (mean ± SD estimated hepatic stores 14 ± 4 mg RE (50 ± 16 µmol RE)). The Panel notes the low hepatic retinol stores of the study population and the short timeframe of the study, which may not have allowed the retinol dose to fully equilibrate with the hepatic pool (see Section 2.4.1.2).

The Panel notes that available data show that the efficiency of storage depends on vitamin A status. Low retinol stores are associated with a reduced efficiency of storage. Data from adult Bangladeshi subjects with liver concentrations ≥ 20 µg retinol/g indicate an average efficiency of storage of ingested retinol of 42 % in the liver. The Panel notes that this would correspond to an efficiency of storage in the whole body of 52 %, assuming that 80 % of retinol body stores are found in the liver in subjects with adequate liver stores.

2.3.5. Metabolism

Retinoic acid is produced from retinol in two oxidative steps. Retinol is first oxidised to retinal, which is further oxidised to retinoic acid.

Two types of enzymes have been implicated in the oxidation of retinol to retinal: the microsomal dehydrogenases of the short-chain dehydrogenases/reductases family of proteins and the cytosolic alcohol dehydrogenases of the medium-chain alcohol dehydrogenases family (Pares et al., 2008). The latter appear to rather play a role as backup enzymes under extreme dietary conditions (Farjo et al., 2011; Napoli, 2012).

The oxidation of retinal to retinoic acid is irreversible. Excessive retinoic acid is catabolised by several cytochrome P450 (CYP) enzymes, giving rise to more water-soluble oxidised and conjugated retinoid forms, which can be more easily excreted (White et al., 1996; Fujii et al., 1997; Ray et al., 1997; White et al., 1997). CYP26A1, CYP26B1 and CYP26C1 appear to be primarily responsible for the degradation of retinoic acid (Pennimpede et al., 2010; Ross and Zolfaghari, 2011; Kedishvili, 2013). With the exception of liver, where CYP26A1 is the predominant form, and lung, where CYP26A1 is slightly more abundant, all other human adult tissues contain higher levels of CYP26B1 transcript (Xi and Yang, 2008; Topletz et al., 2012). Considering that CYP26A1 expression in the liver is very sensitive to retinoic acid levels, the high catalytic efficiency of this low-affinity enzyme would enable CYP26A1 to rapidly bring down excessive levels of retinoic acid. In addition to the three CYP26 enzymes, several other members of other CYP families have been shown to catabolise retinoic acid (Kedishvili, 2013).

Retinal can be converted back to retinol (Kedishvili, 2013), depending on the availability of the substrates and cofactors. The cytosolic aldo-keto reductases and the microsomal short-chain dehydrogenases/reductases have been proposed to catalyse the reduction of retinal back to retinol. This efficient recycling of retinal back to retinol prevents retinal losses through the irreversible pathway to retinoic acid and constitutes a sparing process of retinol stores.

2.3.6. Elimination

2.3.6.1. Catabolic losses

The absolute catabolic rate ($\mu\text{g}/\text{day}$ or $\mu\text{mol}/\text{day}$) and the fractional catabolic rate (% of a defined pool) of retinol are defined as the rate at which retinol is irreversibly utilised each day in absolute or relative amounts, respectively.

The distribution and catabolism of retinol was determined in eight men who received intravenous or oral doses of ^{14}C -labelled retinyl acetate during vitamin A depletion (up to 771 days) and repletion (up to 372 days) (Sauberlich et al., 1974). It took about 26 days for the labelled dose to equilibrate with the total body vitamin pool, which was estimated to range from 315 to 879 mg (1 100–3 070 μmol). A fractional catabolic rate of the total body retinol pool of approximately 0.5 % per day (range 0.3–0.9 %) was determined in these subjects consuming a vitamin A-free diet, deduced from a mean half-life of retinol in the liver of 154 days (range 75–241 days, CV 35 %) during the depletion phase (Sauberlich et al., 1974; Olson, 1987). The absolute utilisation rate of retinol ranged between 1 113 and 2 070 μg (3.9 and 7.2 μmol) per day among subjects at baseline and fell to low levels as depletion progressed (50–180 μg (0.2–0.6 μmol) per day).

The total body retinol content determined by the plasma isotopic ratios of deuterium-labelled retinol was significantly different between groups of four US and Bangladeshi adults (mean \pm SD (range) 295 \pm 13 mg (106–378 mg) (1 030 \pm 45 μmol (370–1 320 μmol)) vs. 286 \pm 315 mg (86–745 mg) (100 \pm 110 μmol (30–260 μmol)), $p = 0.003$) (Haskell et al., 1998). Based on the disappearance kinetics of the fraction of labelled dose in plasma at equilibrium derived from the data of Haskell et al. (1998), Furr et al. (2005) estimated the mean fractional catabolic rate of retinol to be 0.4 % per day (range 0.1–0.7 % per day) in the US subjects and 0.9 % per day (range 0.5–1.2 % per day) in the Bangladeshi subjects. The difference was not statistically significant. It also did not differ from the rate of 0.5 % per day previously determined (Sauberlich et al., 1974).

Based on the same approach, Haskell et al. (2003) estimated a fractional catabolic rate of 2.2 % per day (95 % CI = 1.4–3.0 % per day) in 107 Peruvian children (12–24 months of age) with a total body retinol content (mean \pm SD (range)) estimated as 28 \pm 23 mg (4–112 mg) (97 \pm 81 μmol (16–392 μmol)). According to the authors, the higher fractional catabolic rate in children aged 12–24 months may reflect greater utilisation of the vitamin to support growth, but other factors may have affected the retinol turnover, given that plasma C-reactive protein concentrations were elevated in approximately 50 % of the children. The authors suggested that healthy children (12–24 months of age) may have a fractional catabolic rate lower than 2.2 %.

Applications of model-based compartmental analysis to data from tracer label studies have allowed estimation of parameters of human retinol metabolism, including its catabolic rate (von Reinersdorff et al., 1998; Furr et al., 2005; Cifelli et al., 2008). Such analyses also revealed the important recycling of vitamin A among tissues and plasma before its irreversible utilisation, indicating a sparing process of the vitamin (Reinersdorff et al., 1996; Furr et al., 2005; Cifelli et al., 2008).

Cifelli et al. (2008) investigated retinol kinetics, storage, and catabolic rate through model-based compartmental analysis of data from stable isotope dilution in well-nourished men and women from China (Wang et al., 2004) and the USA (Tang et al., 2003). [$^2\text{H}_8$]-Retinyl acetate (3 mg (8.9 μmol)) was orally administered to US ($n = 12$; 59 \pm 9 years) and Chinese adults ($n = 14$; 54 \pm 4 years) and serum tracer and retinol concentrations were measured from 3 hours to 56 days. Subjects were instructed not to consume vitamin supplements or foods containing large amounts of retinol or β -carotene during the study. Serum retinol concentration was significantly higher in the US group (487 \pm 92 $\mu\text{g}/\text{L}$ (1.70 \pm 0.32 $\mu\text{mol}/\text{L}$)) than in the Chinese group (355 \pm 106 $\mu\text{g}/\text{L}$ (1.24 \pm 0.37 $\mu\text{mol}/\text{L}$), $p < 0.001$) at baseline. Predicted total traced mass (257 \pm 182 vs. 68 \pm 32 mg (898 \pm 637 vs. 237 \pm 109 μmol)), absolute catabolic rate (“disposal rate”) (4.2 \pm 1.7 vs. 1.6 \pm 0.6 mg/day (14.7 \pm 5.87 vs. 5.58 \pm 2.04 $\mu\text{mol}/\text{day}$)) and system residence time (58.8 \pm 28.5 vs.

42.9 ± 14.6 days) were significantly greater in US than in Chinese subjects. In both the US and Chinese participants, absolute retinol catabolic rate was significantly correlated with the traced mass in the extravascular compartment (256 ± 182 and 67 ± 32 mg (892 ± 637 and 233 ± 109 µmol), respectively), with the catabolic rate increasing linearly with increasing stores. The Panel notes that estimated mean daily fractional catabolic rates of 1.6 % (14.7/898) in the US subjects and 2.3 % (5.58/237) in the Chinese subjects would result from the predicted total traced mass and absolute catabolic rate in these two groups, with large inter-individual variability. The Panel notes that the absorption efficiency of retinol (i.e. the portion in compartment 1 that is transferred to compartment 2) estimated by the model is around 65 % in the US subjects. This is likely to underestimate the true absorption, as retinol administered in oil is considered to be almost completely absorbed (Sivakumar and Reddy, 1972) (Section 2.3.1.1). This would lead to an underestimation of the predicted total body pool. Therefore, the fractional catabolic rates derived from these data are likely to overestimate actual fractional catabolic rates.

The Panel notes that the rate of retinol catabolism is related to body stores and that the absolute catabolic rate appears to increase with vitamin A body stores. Overall, retinol catabolism represents a relatively low fraction of the total body pool, owing to the important storage capacity of the body and efficient recycling processes. The Panel notes that available studies were conducted in subjects with a wide range of retinol body stores using different experimental methods and conditions and show substantial variability. The results of the study by Cifelli et al. (2008) indicate that the fractional catabolic rate may be higher than the value of 0.5 % which has usually been considered (Olson, 1987). The Panel notes that the fractional catabolic rate may be influenced by physiological conditions (such as growth, presence of inflammation or other non-identified factors) and that the fractional catabolic rate may be higher in children than in adults, in relation to a higher retinol utilisation for growth needs and, possibly, to relatively lower body stores than in adults.

2.3.6.2. Faecal, breath and urinary losses

The majority of retinol metabolites are excreted in the urine, but they are also excreted in faeces and breath. The percentage of a radioactive dose of ¹⁴C-labelled retinyl acetate recovered in breath, faeces and urine ranged from 18 to 30 %, from 18 to 37 % and from 38 to 60 %, respectively, after 400 days on a vitamin A-deficient diet (Sauberlich et al., 1974). Retinol is metabolised in the liver to numerous products, some of which are conjugated with glucuronic acid or taurine for excretion in bile (Zile et al., 1982; Skare and DeLuca, 1983), and the amount of retinol metabolites excreted in bile increases as the liver retinol exceeds a critical concentration. Excretion of labelled retinol metabolites in bile of rats fed increasing amounts of retinol traced by [³H]-retinyl acetate was constant when hepatic retinol concentrations were low (≤ 32 µg/g (112 nmol/g) and increased rapidly (by eight-fold) as liver retinol concentration increased, up to a plateau at hepatic retinol concentration ≥ 140 µg/g (490 nmol/g) (Hicks et al., 1984). This increased biliary excretion may serve as a protective mechanism for reducing the risk of excess storage of vitamin A.

2.3.6.3. Breast milk

Preformed vitamin A in breast milk primarily occurs as retinyl esters (mainly retinyl palmitate) (Stoltzfus and Underwood, 1995), with a small fraction present as free retinol. Provitamin A carotenoids are also found in breast milk (Canfield et al., 2003). The carotenoid concentration of breast milk is not described in this opinion, as carotenoids are not taken into account in estimating the vitamin A supply in infants, owing to a lack of knowledge on the bioconversion of carotenoids in infants (SCF, 2003; EFSA NDA Panel, 2014b), and secretion of provitamin A carotenoids in breast milk is unlikely to significantly affect the vitamin A status of lactating women.

Preformed vitamin A concentration is higher in colostrum and decreases as lactation progresses (Stoltzfus and Underwood, 1995). It is not related to breast milk fat concentration during the first weeks of lactation (Macias and Schweigert, 2001) but is influenced by maternal vitamin A status (Underwood, 1994b).

Appendix C reports data on retinol⁷ concentration in breast milk from mothers of term infants in Western populations. In a multinational study, Canfield et al. (2003) found mean retinol concentrations between 301 and 352 µg/L in samples of mature milk from Western populations (Australia, Canada, the UK and the USA). Studies on samples taken during the first six months of lactation reported average retinol concentrations in mature milk of 831 µg/L in Germany (Schweigert et al., 2004), 815 µg/L in Turkey (Tokusoglu et al., 2008) and 571 µg/L in Poland (Duda et al., 2009). In a group of Polish lactating women, Kasparova et al. (2012) found decreasing concentrations of retinol in mature breast milk over the course of lactation, from 458 µg/L at 1–2 months post partum to 229 µg/L at 5–6 months post partum and 172 µg/L at 9–12 months post partum.

During the first six months of lactation, the Panel notes that available data indicate that mean total retinol concentrations in mature breast milk of populations from Western countries range between 229 and 831 µg/L. Average values between 450 and 600 µg/L have previously been considered by other committees (DH, 1991; SCF, 1993; Afssa, 2001; IOM, 2001; WHO/FAO, 2004; D-A-CH, 2013; Nordic Council of Ministers, 2014). Based on a mean milk transfer of 0.8 L/day (Butte et al., 2002; FAO/WHO/UNU, 2004; EFSA NDA Panel, 2009) and a concentration of retinol in breast milk of 530 µg/L taken as the midpoint of the range of means (229–831 µg/L), a secretion of 424 µg/day of retinol in breast milk is estimated during the first six months of lactation.

2.3.7. Interaction with other nutrients

Serum retinol concentration was positively associated with serum iron and ferritin concentrations in children (Bloem et al., 1989). Vitamin A deficiency impairs iron mobilisation and vitamin A supplementation improves haemoglobin concentrations (Lynch, 1997). Iron supplementation combined with vitamin A was more effective than iron alone in improving haemoglobin concentrations in anaemic children (Mwanri et al., 2000) and pregnant and lactating women (Suharno et al., 1993; Tanumihardjo et al., 1996; Tanumihardjo, 2002). In a systematic review, vitamin A supplementation during pregnancy was found to reduce anaemia risk (haemoglobin < 11.0 g/dL) among both anaemic and non-anaemic women (Thorne-Lyman and Fawzi, 2012). This is consistent with observational and intervention studies in women and children which showed correlations between anaemia and vitamin A deficiency and the amelioration of anaemia by improving vitamin A status in deficient populations (Radhika et al., 2002; Semba and Bloem, 2002; Al-Mekhlafi et al., 2013). In non-anaemic subjects, a test meal containing 1 000 µg retinol did not enhance iron absorption (Walczyk et al., 2003). Iron deficiency was shown to alter the distribution of retinol and retinyl esters between plasma and liver and to reduce plasma retinol concentrations in rats, despite adequate vitamin A intake and hepatic retinol stores (Amine et al., 1970; Staab et al., 1984; Rosales et al., 1999).

Zinc is important in protein synthesis. In animal models, zinc deficiency affects RBPs and transport of retinol from the liver to the circulation (Terhune and Sandstead, 1972; Smith et al., 1974; Duncan and Hurley, 1978; Baly et al., 1984). In addition, zinc deficiency also reduced the synthesis of rhodopsin in rats (Dorea and Olson, 1986). However, no consistent relationship between zinc and vitamin A status has been established in humans (Christian and West, 1998).

2.3.8. Retinol equivalents

In tissues, blood, milk and food, vitamin A contents are conventionally expressed as RE, with 1 µg RE equal to 1 µg retinol.

The vitamin A activity of provitamin A carotenoids in diets is determined from specific relations between provitamin A carotenoids and retinol to account for the less efficient absorption of carotenoids and their bioconversion to retinol. Conversion factors of 1:6 for β-carotene and 1:12 for other provitamin A carotenoids were initially proposed (FAO/WHO, 1988; SCF, 1993), based on data indicating that 3 µg of dietary β-carotene is equivalent to 1 µg of purified β-carotene in oil and that the

⁷ That is, after saponification to release retinol from retinyl esters.

β -carotene:retinol equivalency ratio of purified β -carotene in oil is approximately 2:1 (Sauberlich et al., 1974). β -Carotene is the most potent retinol precursor of all provitamin A carotenoids (Harrison, 2012). Stoichiometric conversion of 1 mole of β -carotene (with two β -ionone rings) would give rise to 2 moles of retinol (via retinal), whereas conversion of 1 mole of either β -cryptoxanthin or α -carotene (each with only a single β -ionone ring) would give rise to a single mole of retinol. α -Carotene and β -cryptoxanthin show 30–50 % of the provitamin A activity of β -carotene (Bauernfeind, 1972; van Vliet et al., 1996).

In 2001, the Institute of Medicine (IOM) revised these factors considering new data (IOM, 2001): (1) absorption of β -carotene from a mixed vegetable diet had been reported to be 14 % compared with absorption of β -carotene in oil (van het Hof et al., 1999); (2) absorption from green leafy vegetables appeared to be lower than absorption from fruits (de Pee et al., 1998); and (3) a low proportion of β -carotene was consumed from fruits compared with vegetables in the USA. Retinol activity equivalency⁸ ratios of 1:12 for β -carotene and 1:24 for other provitamin A carotenoids were proposed. Considering the data from van het Hof et al. (1999), WHO/FAO (2004) also proposed revised equivalency factors of 1:14 for β -carotene and 1:28 for other provitamin A carotenoids from usual vegetable diets, with possible adjustment depending on the proportion of green leafy vegetables or fruits in the diet. West et al. (2002) suggested that these revised conversion factors might still be too high, especially for populations living in developing countries. Two recent reviews (Haskell, 2012; Van Loo-Bouwman et al., 2014) provide an overview of experimental data on β -carotene:retinol equivalency ratios published until early 2011 and mid-2013, respectively.

Reviews of data on the bioavailability of β -carotene from plant sources in humans (Haskell, 2012; Van Loo-Bouwman et al., 2014) reported absorption ranging from 5 to 65 % and retinol equivalency ratios for β -carotene ranging from 3.8:1 to 28:1 by weight. In line with de Pee et al. (1998), there was further indication that β -carotene from fruits is better converted than β -carotene from green leafy vegetables (Khan et al., 2007). For pure β -carotene diluted in oil, equivalency ratios from 2:1 to 55:1 were reported, with most values being between 2:1 and 4:1. The data collected by Haskell (2012) indicate that the efficiency of conversion of β -carotene from oil might be increased in subjects with “low or marginal” vitamin A status compared with subjects with “adequate” vitamin A status, while it appears to decrease with increasing dose of β -carotene. Overall, there is high variability in retinol equivalency ratios, which might originate from either host-related factors (genetics, age, sex, nutritional status, digestive dysfunctions and illness) or food-related factors (food composition, food matrix) (de Pee and Bloem, 2007; Tanumihardjo et al., 2010; Haskell, 2012). A study in eight healthy free-living adults who received an oral tracer dose of [¹⁴C]- β -carotene also confirms that β -carotene conversion is highly variable (Ho et al., 2009).

Few results are available on the rate of absorption of β -carotene and its bioequivalence to retinol in children (Van Loo-Bouwman et al., 2014). By measuring the plasma ratio of retinol formed from labelled β -carotene compared with a reference dose of labelled retinol, van Lieshout et al. (2001) estimated that the amount of β -carotene in oil required to form 1 μ g retinol was 2.4 μ g (95 % CI = 2.1–2.7 μ g) in 36 Indonesian children aged 8–11 years. In a study in 68 Chinese children (6–8 years) using labelled retinyl acetate as a reference, the mean (\pm SE) conversion factors of pure β -carotene, β -carotene from Golden Rice and β -carotene from spinach to retinol were 2.0 ± 0.9 , 2.3 ± 0.8 and 7.5 ± 0.8 to 1, respectively (Tang et al., 2012). Ribaya-Mercado et al. showed a significant improvement in vitamin A status, as assessed by the deuterated retinol dilution method, in Filipino schoolchildren receiving controlled diets rich in provitamin A carotenoids from fruit and vegetable sources, but these studies do not allow the estimation of provitamin A carotenoid/retinol equivalency ratios (Ribaya-Mercado et al., 2000; Ribaya-Mercado et al., 2007).

⁸ Retinol Activity Equivalents (RAEs) were defined by IOM (2001) assuming that, after absorption of β -carotene, whether from oil or from food, the metabolism of the molecule is similar and that the retinol equivalency ratio of β -carotene in oil is 2:1. To take into account the lower absorption efficiency of β -carotene from food compared with that from oil, the vitamin A activity of β -carotene from food can be derived by multiplying by 2:1. For example, it was estimated that 6 μ g of β -carotene from a mixed diet is nutritionally equivalent to 1 μ g of β -carotene in oil. Therefore, the retinol activity equivalency (μ g RAE) ratio for β -carotene from food was estimated to be 12:1 ($6 \times 2:1$).

The Panel notes the high variability in the β -carotene/retinol equivalency ratios estimated from these studies, depending on the food matrix, the subjects' vitamin A status and the dose administered. This results in large uncertainties in establishing equivalency ratios from the whole diet of large populations. The Panel considers that current evidence is insufficient to support a change from the conversion factors proposed by the SCF for European populations, namely 1 μg RE equals 1 μg of retinol, 6 μg of β -carotene and 12 μg of other carotenoids with provitamin A activity.

2.4. Biomarkers

2.4.1. Total body retinol content and liver retinol concentration

Vitamin A status can best be expressed in terms of total body pool of retinol (i.e. as free retinol and retinyl esters) or, alternatively, in terms of liver concentration of the vitamin (Olson, 1987). Hepatic concentration is considered as a marker of vitamin A status because 70–90 % of the retinol in the body is stored in the liver in healthy individuals, while this percentage is considered to decrease to 50 % or below in severely deficient individuals (Rietz et al., 1974; Bausch and Rietz, 1977; Olson, 1987) (Section 2.3.4.1).

Olson (1987) has proposed a minimum concentration of 20 μg retinol/g liver (0.07 $\mu\text{mol/g}$) (i.e. as free retinol and retinyl esters) as a criterion to define adequate vitamin A status, based on the following considerations: (1) no clinical signs of deficiency have been noted in individuals with this or a higher liver concentration; (2) at this concentration and above, the liver is capable of maintaining a steady-state plasma retinol concentration, as determined by the relative dose–response test in rats (Loerch et al., 1979) and humans (Amedee-Manesme et al., 1987); (3) biliary excretion of retinol increases significantly when liver stores rise significantly above this concentration in rats (Hicks et al., 1984), which is suggested to serve as a regulatory mechanism of vitamin A storage; and (4) this concentration was calculated to be sufficient to protect an adult ingesting a diet free of vitamin A from a deficiency state for approximately four months as well as to meet vitamin A needs during shorter periods of stress (e.g. infection).

This value has commonly been used as a reference point to define adequate vitamin A status in the scientific literature (Olson, 1987), as well as to derive vitamin A requirements (SCF, 1993; IOM, 2001; WHO/FAO, 2004). The Panel considers that a concentration of ≥ 20 μg retinol/g liver (0.07 $\mu\text{mol/g}$) can be considered to reflect an adequate vitamin A status.

2.4.1.1. Direct measurement

Liver retinol concentrations have been directly determined by post-mortem liver analysis and analysis of biopsy samples. Post-mortem liver analysis indicated concentrations of retinol ranging from 10 to 1 807 $\mu\text{g/g}$ liver (0.03 to 6.3 $\mu\text{mol/g}$) in Western countries (Hoppner et al., 1969; Underwood et al., 1970; Raica et al., 1972; Mitchell et al., 1973; Money, 1978; Huque, 1982; Schindler et al., 1988). Mean and median retinol concentrations were 252 $\mu\text{g/g}$ (0.9 $\mu\text{mol/g}$) and 198 $\mu\text{g/g}$ (0.7 $\mu\text{mol/g}$) (range 0–1 201 $\mu\text{g/g}$ (0–4.2 $\mu\text{mol/g}$) in post-mortem analysis of the liver of 364 British males and females (aged 0 to >90 years) (Huque, 1982). Mean (\pm SD) and median retinol concentrations of 597 \pm 397 $\mu\text{g/g}$ (2.1 \pm 1.4 $\mu\text{mol/g}$) and 506 $\mu\text{g/g}$ (1.8 $\mu\text{mol/g}$) (range 36–1 807 $\mu\text{g/g}$ (0.1–6.3 $\mu\text{mol/g}$)) were found in post-mortem analysis of the liver of 77 men and women (mean age 56 years) in Germany (Schindler et al., 1988). Analysis of liver biopsy samples performed in low-income countries showed hepatic concentrations of 17–141 $\mu\text{g/g}$ (0.1–0.5 $\mu\text{mol/g}$) (Suthutvoravoot and Olson, 1974; Abedin et al., 1976; Olson, 1979; Flores and de Araujo, 1984; Furr et al., 1989; Haskell et al., 1997). However, post-mortem liver analysis and liver biopsies are not feasible in population-based studies as primary status indicators for obvious reasons.

2.4.1.2. Indirect measurement by stable isotope dilution methods

Total body content and liver retinol concentration can be estimated indirectly by stable isotope dilution approaches (Haskell et al., 2005; IAEA, 2008). After oral administration of a small dose of retinol tracer labelled with stable deuterium (^2H) or carbon (^{13}C), the dilution of tracer in plasma is measured

when the labelled dose has mixed with endogenous stores and equilibrium is reached (14–20 days after administration). Total body exchangeable retinol pool can be derived from a mass balance equation, correcting for the efficiency of absorption and storage of retinol and its fractional catabolic rate (Furr et al., 1989; Furr et al., 2005; IAEA, 2008).

In the deuterated retinol dilution (DRD) technique, the retinol pool is calculated from an equation developed by Furr et al. (1989),⁹ considering efficiency of absorption and storage of retinol, its catabolic rate and inequality of the plasma to liver ratio of labelled to non-labelled retinol. The absorption and storage efficiency factor is usually assumed to be 50 % based on data from Bausch and Rietz (1977) (see Section 2.3.4.3). To adjust for the catabolism of the labelled dose during the equilibration period, a fractional catabolic rate of 0.5 % is typically considered, derived from the half-life of retinol turnover in adults (Saubert et al., 1974) (see Section 2.3.6.1). To account for the fact that unlabelled retinol is continuously consumed in the diet and newly absorbed retinol contributes preferentially to the plasma pool, another factor is applied to correct for the difference in specific activity in liver compared with plasma. A value of 0.65 is usually assumed, derived from the ratio observed in rats (Hicks et al., 1984). This factor is not needed if no retinol, or as little as possible, is consumed during the equilibration period.

In the [¹³C₂]-retinol isotope dilution ([¹³C₂]-RID) test, the dose of tracer administered is smaller than in the DRD technique, which reduces the degree to which the dose perturbs the endogenous retinol pool (Furr et al., 2005). For this test, a dose absorption of 90–100 % is assumed and there is no correction for the differences in distribution of the tracer between liver and serum (Valentine, 2013).

For both techniques, hepatic stores can be further determined by considering that the amount of retinol stored in the liver is positively correlated with the size of the total body pool. Between 40 and 90 % of the total body retinol pool is assumed to be stored in the liver, depending on the vitamin A status of the subjects (Rietz et al., 1974; Bausch and Rietz, 1977) (Section 2.3.4.1).

Based on data from 10 adult subjects in the USA, the correlation coefficient between liver retinol concentrations calculated from the DRD method (range 19–321 µg/g liver (0.065–1.12 µmol/g)) and directly measured in liver biopsies (range 14–160 µg/g liver (0.049–0.56 µmol/g)) was 0.88, and the Spearman's rank correlation coefficient was 0.95 ($p < 0.002$) (Furr et al., 1989). Based on data from 31 Bangladeshi surgery patients, Haskell et al. (1997) found good agreement between mean hepatic stores of retinol estimated by the DRD technique (32 ± 21 mg (0.110 ± 0.072 mmol)) and by liver biopsy (29 ± 19 mg (0.100 ± 0.067 mmol)), assuming that liver weight is 2.4 % of body weight. A significant linear relationship was found between the two techniques ($r = 0.75$, $p < 0.0001$). However, a wide prediction interval was observed for estimates of hepatic retinol stores for individual subjects.

Liver and total body retinol stores assessed by stable isotope dilution methods have been shown to be well correlated with measures of habitual vitamin A intake in cross-sectional studies over a wide range of intakes (Pearson correlation coefficients around 0.4) (Ribaya-Mercado et al., 2004; Valentine et al., 2013) and to respond to vitamin A supplementation in intervention studies lasting a couple of weeks (Haskell et al., 1999; Ribaya-Mercado et al., 1999; Haskell et al., 2011).

⁹ Total body exchangeable vitamin A pool = $F \times \text{dose} \times [S \times a \times ((1/D:H) - 1)]$, where:

- F is a factor related to the efficiency of absorption and storage of the orally administered dose;
- dose is the amount of isotope orally administered (mmol);
- the factor S corrects for the inequality of the plasma to liver ratio of labelled to non-labelled retinol; this correction is not needed if subjects consume as little vitamin A as possible after administering the oral dose, while the isotope is mixing with exchangeable vitamin A pools;
- the factor a corrects for irreversible loss of labelled vitamin A during the equilibration period;
- D:H is the isotopic ratio of labelled to non-labelled retinol in plasma;
- and -1 corrects for the contribution of the dose to the total body vitamin A reserve (this term is omitted when the mass of the labelled vitamin A is small compared with the mass of total body vitamin A).

The Panel notes that there are uncertainties inherent in stable isotope dilution methods owing to the assumptions required in the calculations. Human data on the parameters used are limited, so that inter-individual variability and the influence of factors such as age are not well characterised (IAEA, 2008). The methods also assume that the fractional catabolic rate is independent of the size of the stores of retinol, which is unlikely, as indicated by data in rats (Green and Green, 1994) and humans (Sauberlich et al., 1974; Cifelli et al., 2008) (Section 2.3.6.1). Despite these limitations, they have the advantage that they enable a quantitative estimation of retinol stores. The Panel notes that these methods provide good estimates at group level, but lack precision for their determination at individual level, owing to the large inter-individual variability of the factors used in the equation.

2.4.1.3. Relative dose–response

The relative dose–response (RDR) is an indirect measurement of hepatic retinol stores. In conditions of vitamin A deficiency, RBP that is not bound to retinol (apo-RBP) accumulates in the liver, but is rapidly released from the liver into the circulation after administration of a dose of retinol or retinyl esters (Muto et al., 1972; Smith et al., 1973; Loerch et al., 1979).

In this test, after an oral dose of retinol, the relative excess of apo-RBP in the liver binds to retinol and the resulting holo-RBP (RBP bound with retinol), coupled with transthyretin, is released into the circulation. Two blood samples are collected, at baseline and five hours after dosing, and the RDR value is calculated as follows:

$$\text{RDR (in \%)} = \left[\frac{\text{serum retinol concentration at five hours post-dosing} - \text{serum retinol concentration at baseline}}{\text{serum retinol concentration at five hours post-dosing}} \right] \times 100.$$

Alternative methods have been proposed, e.g. by measuring serum RBP instead of serum retinol concentration (Fujita et al., 2009), or by administering the 3,4-didehydroretinyl ester analogue instead of retinol as the test dose (modified RDR) (Tanumihardjo, 1993).

The RDR test is considered a valid test to determine inadequate vitamin A status. A large positive response to the dose, i.e. an RDR value > 20 %, is indicative of vitamin A deficiency, whereas a value < 20 % is considered to reflect hepatic concentrations equivalent to or above 20 µg retinol/g (0.07 µmol/g) (Tanumihardjo, 1993; WHO, 1996; Tanumihardjo, 2011). However, the synthesis of RBP also depends on the adequacy of energy intake and of other nutrients such as zinc and protein. In addition, plasma retinol concentration and, consequently, the RDR test are insensitive across a wide range of liver concentrations above 20 µg retinol/g (0.07 µmol/g) (Solomons et al., 1990).

The Panel considers that the RDR represents a good marker of inadequate vitamin A status, but its sensitivity is limited to liver concentrations below 20 µg retinol/g (0.07 µmol/g).

2.4.2. Plasma/serum retinol concentration

In the usual range, plasma/serum retinol concentration is neither related to observed habitual vitamin A intake from either dietary preformed vitamin A or provitamin A carotenoid sources nor responsive to supplement use (IOM, 2001; Tanumihardjo, 2011).

The concentration of plasma/serum retinol is under tight homeostatic control (Olson, 1984). The relationships between plasma/serum retinol and total body retinol content or liver retinol concentration are not linear. Plasma/serum retinol concentrations reflect liver retinol stores only when they are severely depleted (< 20 µg retinol/g liver (< 0.07 µmol/g)) or very high (> 300 µg/g liver (1.05 µmol/g)) (WHO, 2011). A plasma/serum retinol concentration below 200 µg/L (0.7 µmol/L) is considered to reflect vitamin A inadequacy for population assessment (Sommer, 1982; Olson, 1987; Flores, 1993; Underwood, 1994a; WHO, 2011). The prevalence of values below 200 µg/L (0.7 µmol/L) is a generally accepted population cut-off for preschool-age children to indicate risk of inadequate vitamin A status (WHO, 1996, 2011), whilst values above 300 µg/L (1.05 µmol/L) indicate

an adequate status related to the absence of clinical signs of deficiency (Pilch, 1987; Flores et al., 1991).

A low plasma/serum retinol concentration may also originate from an inadequate supply of dietary protein, energy, or zinc, which are required for synthesis of RBP, or may be caused by infection and inflammation resulting in decreases in the concentrations of some acute phase proteins, such as RBP and transthyretin (IOM, 2001; Tanumihardjo, 2011). Infection can lower mean plasma/serum retinol concentration by as much as 25 %, independently of vitamin A intake (Filteau et al., 1993; Christian et al., 1998b).

The Panel notes that the specificity of plasma/serum retinol concentration is affected by a number of factors unrelated to vitamin A status, including infection and inflammation, which make the interpretation of this biomarker difficult. In addition, plasma/serum retinol concentrations are maintained nearly constant over a wide range of vitamin A intakes.

2.4.3. Markers of visual function

Xerophthalmia is the most specific vitamin A deficiency disorder (Section 2.2.2.1). It encompasses the clinical spectrum of ocular manifestations of vitamin A deficiency, from milder stages of night blindness and Bitot's spots, to potentially blinding stages of corneal xerosis, ulceration and necrosis (WHO, 2009). The prevalence of xerophthalmia is considered a population indicator of vitamin A deficiency (WHO, 2009; Tanumihardjo, 2011).

2.4.3.1. Night blindness

The rhodopsin molecule of the rods in the retina contains 11-*cis* retinal (Section 2.2.1). Without an adequate supply of vitamin A to the retina, the function of the rods in dim light situations is affected, resulting in abnormal dark adaptation, i.e. night blindness (Carney and Russell, 1980).

Numerous tests have been used to assess the presence of night blindness (WHO, 2012). The most common method used at population level involves subjective reports on current or past night blindness status. Objective measures have also been developed based on dark adaptation or the scopic response to various light stimuli after dark adaptation. They include dark adaptometry, the pupillary response test and the night vision threshold test.

Measures of night blindness and dark adaptometry are sensitive markers of vitamin A status at the lower end of the status continuum (liver concentration < 20 µg retinol/g (0.07 µmol/g)) (Tanumihardjo, 2011). Epidemiological evidence suggests that host resistance to infection is impaired prior to clinical onset of night blindness and laboratory animals fed a vitamin A-deficient diet maintain ocular concentrations of vitamin A despite a significant reduction in liver retinol concentration (IOM, 2001).

In addition, zinc deficiency and severe protein deficiency also may affect dark adaptation responses (Morrison et al., 1978; Bankson et al., 1989).

2.4.3.2. Conjunctival impression cytology

Vitamin A deficiency leads to early keratinising metaplasia (Bitot's spots) and losses of mucin-secreting goblet cells on the bulbar surface of the conjunctiva of the eye (IOM, 2001). Cells can be counted and evaluated by microscopic examination of a filter paper impression from the surface of the eye and staining with haematoxylin and eosin (Tanumihardjo, 2011). However, there have been concerns about the performance of this method to assess vitamin A deficiency when compared with biochemical markers (e.g. serum retinol or RDR) (Amedee-Manesme et al., 1988; Gadomski et al., 1989; Rahman et al., 1995; Sommer and West, 1996). This technique was used in the 1990s, but, because of its limitations, has not been widely adopted (Tanumihardjo, 2011).

2.4.3.3. Conclusion on markers of visual function

The Panel notes that markers of visual function have been used for population evaluation of vitamin A status or to assess intervention efficacy. However, these methods are rather qualitative and their sensitivity is limited to situations of vitamin A deficiency (Tanumihardjo, 2011).

2.4.4. Conclusion on biomarkers

The Panel notes that plasma/serum retinol concentration is under tight homeostatic control and does not reflect vitamin A status until body stores are very low (or very high). In contrast, measures of total body content or liver concentration by stable isotope dilution methods have shown good correlation with habitual vitamin A intake, over a wide range of intakes.

As reviewed by Tanumihardjo (2011), the sensitivity of markers of visual function is limited to situations of vitamin A deficiency. Likewise, the RDR is a useful marker of inadequate status only. In contrast, stable isotope dilution methods give a quantitative estimate of liver retinol concentrations over a broad range, from vitamin A deficiency to toxicity.

The Panel considers that measurements of the total body retinol content or liver retinol concentration are the most specific and sensitive markers of vitamin A status. Liver concentration $< 20 \mu\text{g}$ retinol/g ($0.07 \mu\text{mol/g}$) (i.e. as free retinol and retinyl esters) can be used as an indicator of vitamin A deficiency, while concentrations above this value are considered to maintain adequate plasma retinol concentrations, prevent clinical signs of deficiency and reflect adequate vitamin A status.

2.5. Effects of genotypes

In recent years, large subsets of molecular components of the metabolism of retinoids have been identified (D'Ambrosio et al., 2011). Mutations in the corresponding genes can cause various diseases, including retinitis pigmentosa and Stargardt disease, that lead to severe vision impairment and often blindness (Palczewski, 2010). Moreover, mutations in these genes can cause Matthew–Wood syndrome, a fatal disease which is associated with anophthalmia, pulmonary and cardiac malfunctions and severe mental retardation (Blaner, 2007).

Many proteins participate in the processes involved in the intestinal metabolism of retinol and carotenoids. Given the important role of these proteins in the absorption of dietary carotenoids, their conversion to retinol, and the incorporation of both carotenoids and retinyl esters into chylomicrons, it is not surprising that polymorphisms in these genes affect carotenoid transport and metabolism (Erlanson and Borgstrom, 1968; von Lintig, 2010). Single nucleotide polymorphisms in SR-B1 (Borel et al., 2007) and in BCMO1 (Ferrucci et al., 2009; Leung et al., 2009) have been associated with alterations in the metabolism of carotenoids and retinoids in humans. In humans, a heterozygotic mutation in BCMO1 was described with evidence of both elevated plasma β -carotene concentration and low plasma retinol concentration (Lindqvist et al., 2007).

Mutations in the retinal pigment epithelium-specific 65 kDa protein (RPE65) in humans result in chromophore deficiency and blindness (Marlhens et al., 1997).

The Panel considers that genotype probably induces inter-individual differences in vitamin A requirement but present knowledge is limited and cannot be used for setting DRVs for vitamin A.

3. Dietary sources and intake data

3.1. Dietary sources

Foods rich in retinol include offal and meat, butter, retinol-enriched margarine, dairy products and eggs, while foods rich in provitamin A carotenoids, in particular β -carotene, include vegetables and fruits, such as sweet potatoes, carrots, pumpkins, dark green leafy vegetables, sweet red peppers, mangoes and melons (FSA, 2002; Anses/CIQUAL, 2012).

Currently, vitamin A (as retinol, retinyl acetate, retinyl palmitate and β -carotene) may be added to foods¹⁰ and food supplements.¹¹ The vitamin A content of infant and follow-on formulae¹² and of processed cereal-based foods and baby foods for infants and young children¹³ is regulated.

3.2. Dietary intake

EFSA estimated dietary intake of vitamin A from food consumption data from the EFSA Comprehensive Food Consumption Database (EFSA, 2011b), classified according to the food classification and description system FoodEx2 (EFSA, 2011a). Data from 12 dietary surveys in nine EU countries were used. The countries included were Finland, France, Germany, Ireland, Italy, Latvia, the Netherlands, Sweden and the UK. The data covered all age groups from children to adults (Appendix D).

Nutrient composition data for vitamin A were derived from the EFSA Nutrient Composition Database (Roe et al., 2013). The vitamin A content of foods, expressed as RE, was calculated by considering that 1 μg RE equals 1 μg retinol and 6 μg β -carotene. Other provitamin A carotenoids (i.e. α -carotene and β -cryptoxanthin) were not taken into account because of the limited availability of data concerning these compounds in the database.

Food composition information from Finland, Germany, Italy, the Netherlands and the UK was used to calculate vitamin A intake in these countries, assuming that the best intake estimate would be obtained when both the consumption data and the composition data are from the same country. For vitamin A intake estimates of Ireland and Latvia, food composition data from the UK and Germany, respectively, were used, because no specific composition data from these countries were available. EFSA intake estimates are based on consumption of foods, either fortified or not (i.e. without dietary supplements).

Average vitamin A intake ranged between 409 and 651 μg RE/day in children aged 1 to < 3 years, between 607 and 889 μg RE/day in children aged 3 to < 10 years, between 597 and 1 078 μg RE/day in children aged 10 to < 18 years, and between 816 and 1 498 μg RE/day in adults. Average daily intakes were in most cases slightly higher in males (Appendix E) than in females (Appendix F), mainly owing to the larger quantities of food consumed per day.

Among children aged 1 to < 3 years, food products for the young population, vegetables and vegetable products, milk and dairy products contributed significantly to vitamin A intake. In the older age groups, in addition to the vegetable and vegetable products and milk and dairy products, meat and meat products and animal and vegetable fats also contributed to vitamin A intake (Appendices G and H). Differences in the main contributors to vitamin A intake between sexes were minor.

When available, EFSA vitamin A intake estimates were compared with published intake estimates from the same national surveys. EFSA intake estimates were found to differ by 1–10 % from the published results of the EsKiMo and VELS surveys in Germany (Kersting and Clausen, 2003; Mensink et al., 2007), the National Diet and Nutrition Survey in the UK (Bates et al., 2012) and the National Adult Nutrition Survey in Ireland (IUNA, 2011). Differences were found to be higher, up to 24 %, compared with published results from the INCA 2 survey in France (Afssa, 2009) and the third INRAN-SCAI survey in Italy (Sette et al., 2011). Comparisons were not possible for Finland (Helldán et al., 2013), Sweden (Amcoff et al., 2012) and the Netherlands (van Rossum et al., 2011) owing to the use of different conversion factors for provitamin A carotenoids for calculating vitamin A content of foods. Uncertainties in the estimates may be caused by differences in disaggregating data for

¹⁰ Regulation (EC) No 1925/2006 of the European Parliament and of the Council of 20 December 2006 on the addition of vitamins and minerals and of certain other substances to foods. OJ L 404, 30.12.2006, p. 26.

¹¹ Directive 2002/46/EC of the European Parliament and of the Council of 10 June 2002 on the approximation of the laws of the Member States relating to food supplements. OJ L 183, 12.7.2002, p. 51.

¹² Commission Directive 2006/141/EC of 22 December 2006 on infant formulae and follow-on formulae and amending Directive 1999/21/EC. OJ L 401, 30.12.2006, p. 1.

¹³ Commission Directive 2006/125/EC of 5 December 2006 on processed cereal-based foods and baby foods for infants and young children. OJ L 339, 06.12.2006, p. 16–35.

composite dishes before intake estimations; inaccuracies in mapping food consumption data according to the FoodEx2 classification; analytical errors or errors in estimating vitamin A content of foods in the food composition tables; the use of borrowed vitamin A values from other countries; or the replacement of missing vitamin A contents for some foods by values for similar foods or food groups in the vitamin A intake estimation process. It is not possible to conclude which of these intake estimates would be closer to the actual vitamin A intake.

4. Overview of Dietary Reference Values and recommendations

4.1. Adults

In their revision of the Nordic Nutrition Recommendations (NNR), the Nordic countries decided to maintain their earlier recommendations of 900 µg RE/day for men and 700 µg RE/day for women (Nordic Council of Ministers, 2014), which was based on the approach adopted by IOM (2001). A study in men using the DRD technique to estimate vitamin A requirement was noted (Haskell et al., 2011), but it was considered that more studies on the variation in the AR were needed before a change in the current recommendations could be proposed.

D-A-CH (2013) derived an AR for men of 600 µg RE/day, which was reported to have been determined experimentally. Using a CV of 30 %, a recommended intake of 1 000 µg RE/day for men was proposed. The recommended intake for women was set at 800 µg RE/day, i.e. 20 % less than that for men, considering that their average plasma concentration is lower (Heseker et al., 1994).

At the FAO/WHO Expert consultation of 1998 (WHO/FAO, 2004), experts maintained the approach that had been proposed previously (FAO/WHO, 1988). The mean requirement¹⁴ was defined as the minimum daily intake of vitamin A required to prevent xerophthalmia in the absence of clinical or subclinical infection. A mean requirement of 4–5 µg/kg body weight per day was estimated from the depletion–repletion study by Sauberlich et al. (1974). Vitamin A mean requirements of 300 µg RE/day for men and 270 µg RE/day for women were proposed. The “safe level of intake” was defined as the average continuing intake of vitamin A required to permit vitamin A-dependent functions and to maintain an acceptable total body store of the vitamin. This store helps to offset periods of low intake or increased need resulting from infection and other forms of stress. Recommended Safe Intakes of 500 µg RE/day for women aged 19–65 years, 600 µg RE/day for women aged > 65 years and 600 µg RE/day for men were set. These were calculated by estimating the average dietary intake of retinol needed to replace the endogenous stores that are lost,¹⁵ following the approach proposed by Olson (1987), and considering a CV of 20 %. The CV was estimated from data on the half-life of vitamin A reported by Sauberlich et al. (1974). Equivalency factors of 1:14 for β-carotene and 1:28 for other provitamin A carotenoids from usual vegetable diets were recommended (van het Hof et al., 1999), which may be adjusted depending on the proportion of green leafy vegetables or fruits in the diet.

IOM (2001) estimated the requirement for vitamin A based on the assurance of adequate stores of vitamin A. A minimum acceptable liver vitamin A concentration of 20 µg/g (0.07 µmol/g) was considered. At this concentration, no clinical signs of deficiency are observed, adequate plasma retinol concentrations are maintained (Loerch et al., 1979), induced biliary excretion of vitamin A is observed (Hicks et al., 1984) and protection against vitamin A deficiency is ensured for approximately four months while consuming a vitamin A-deficient diet. The Estimated Average Requirement (EAR) was calculated by multiplying the percentage of body vitamin A stores lost per day when ingesting a vitamin A-free diet (0.5 %), the minimum acceptable liver retinol concentration (20 µg/g), the liver weight:body weight ratio (1:33), the reference weight for a specific age group and sex (61 and 76 kg for women and men, respectively), the ratio of total body:liver vitamin A stores (10:9) and the efficiency of storage of ingested vitamin A (40 %) (Olson, 1987). This resulted in an EAR of 627 µg Retinol Activity Equivalent (RAE)/day for men and 503 µg RAE/day for women. A CV of 20 % was

¹⁴ Previously defined as “basal requirement” by FAO/WHO (1988).

¹⁵ Previously defined as “mean normative storage requirement” by FAO/WHO (1988).

used to derive the Recommended Dietary Allowance (RDA) based on the estimated half-life of vitamin A. RDAs of 900 µg RAE/day for men and 700 µg RAE/day for women were set. The IOM revised conversion factors of carotenoids and retinol to account for data suggesting a lower absorption of provitamin A carotenoids (de Pee et al., 1998; Parker et al., 1999; van het Hof et al., 1999). Retinol activity equivalency factors of 12:1 for dietary β-carotene and 24:1 for other dietary provitamin A carotenoids were proposed (Section 2.3.8).

Afssa (2001) considered a vitamin A requirement of 600 µg RE/day, based on data from the depletion–repletion study by Hume and Krebs (1949) and the radioisotope study by Sauberlich et al. (1974). Given the small number of subjects involved in these studies, an individual variation of 30 % was considered and a PRI of 800 µg RE/day for men was proposed. For women, the value was extrapolated from the value for men on the basis of energy requirements and set at 600 µg RE/day. Afssa recommended 350 µg RE/day to be provided by β-carotene (2.1 mg/day). The vitamin A activity of carotenoids in the diet was expressed in RE based on conversion factors of 6:1 for dietary vitamin A:β-carotene and 12:1 for vitamin A:other dietary provitamin A carotenoids. Because older adults may be at particular risk for hypervitaminosis A due to protein deficiency or renal failure, Afssa proposed to set the PRI at 700 µg RE/day for men and 600 µg RE/day for women over 75 years (Ward, 1996).

The SCF (1993) considered the approach proposed by Olson (1987), using a liver concentration of 20 µg retinol/g (0.07 µmol/g) as a criterion for vitamin A sufficiency. The mean dietary intake needed to maintain this concentration was calculated assuming that the liver store represents 90 % of the total body vitamin A pool and the efficiency of storage in the liver is 50 %. Based on studies with radioactive vitamin A, a mean fractional catabolic rate of 0.5 % was considered. This results in a mean daily dietary intake of 6.7 µg RE/kg body weight, corresponding to an AR of 500 µg RE/day for men and 400 µg RE/day for women. A CV of 20 % was considered from the rates of depletion observed experimentally. The PRI was set at 700 µg RE/day for men and 600 µg RE/day for women. Conversion factors of 6:1 for dietary vitamin A:β-carotene and 12:1 for vitamin A:other dietary provitamin A carotenoids were recommended.

The Netherlands Food and Nutrition Council (1992) identified a minimum requirement of vitamin A for adults of 600 µg RE/day, which was observed to be sufficient to prevent deficiency symptoms such as electroretinographic anomalies and changes in the eyes and the skin, and to maintain plasma retinol concentration at a minimum of 0.7 µmol/L (Sauberlich et al., 1974). An Adequate intake (AI) of 1 000 µg RE/day for men and 800 µg RE/day for women was proposed.

The UK Committee on Medical Aspects of Food Policy (COMA) (DH, 1991) referred to the approach proposed by FAO/WHO (1988), which based recommendations on the maintenance of an adequate body pool size, considering the concentration of vitamin A in the liver. The PRIs were set at 700 µg RE/day for men and 600 µg RE/day for women.

Table 1: Overview of Dietary Reference Values for vitamin A for adults

	NCM (2014)	D-A-CH (2013)	WHO/FAO (2004)	Afssa (2001)	IOM (2001)	SCF (1993)	NL (1992)	DH (1991)
Age (years)	≥ 19	≥ 19	19–65	≥ 19	≥ 19	All	≥ 19	Adults
PRI Men (µg RE/day)	900	1 000	600 ^(a)	800	900 ^(b)	700	1 000 ^(c)	700
PRI Women (µg RE/day)	700	800	500 ^(a)	600	700 ^(b)	600	800 ^(c)	600
Age (years)			> 65	≥ 75				
PRI Men (µg RE/day)			600 ^(a)	700				
PRI Women (µg RE/day)			600 ^(a)	600				

NCM, Nordic Council of Ministers; NL, Netherlands Foods and Nutrition Council; PRI, Population Reference Intake; RE, Retinol Equivalent; RAE, Retinol Activity Equivalent.

(a): Recommended Safe Intake.

- (b): Expressed in μg RAE/day.
(c): Adequate Intake.

4.2. Infants and children

The Nordic countries maintained their earlier approach and extrapolated recommended intakes for children and adolescents from those for adults by using metabolic body weight (body weight to the power of 0.75) and growth factors (Nordic Council of Ministers, 2014).

From observed intakes of breast-fed infants in developing countries with no signs of deficiency and normal growth, WHO/FAO (2004) estimated a requirement of 180 μg RE/day for infants aged 0–6 months and increased it to 190 μg RE/day for infants aged 7–12 months. Considering vitamin A intakes from breast milk in well-nourished communities, a Recommended Safe Intake of 375 μg RE/day was proposed in early infancy ($1.75 \mu\text{mol/L} \times 0.75 \text{ L/day}$) and increased to 400 μg RE/day for infants aged 7–12 months, taking into consideration that vitamin A-deficient populations are at increased risk of death from six months onwards. The requirement and Recommended Safe Intake for preschool children were derived from the values set in late infancy (i.e. 20 and 39 μg RE/kg body weight per day) and estimated to be in the range of 200–400 μg RE/day. Such values were supported by intakes observed to relieve signs of deficiency and reduce the risk of mortality in Indian children (Rahmathullah et al., 1990) and maintain serum retinol concentrations of 0.7 $\mu\text{mol/L}$ in American preschool children (Ballew et al., 2001). Recommendations for older children were derived from adult values.

IOM (2001) proposed an AI of 500 μg RAE/day for infants aged 7–12 months, considering that the extrapolation from the AI set for infants aged 0–6 months fed breast milk resulted in an estimate of 483 μg RAE/day, and that the estimation of total intakes based on the calculated intake from human milk ($485 \mu\text{g/L} \times 0.6 \text{ L/day} = 291 \mu\text{g/day}$) and observed intake from complementary foods (244 $\mu\text{g/day}$, $n = 44$, Third National Health and Nutrition Examination Survey) resulted in an estimate of 535 μg RAE/day. For children and adolescents, no data were available to estimate an average requirement. EARs were extrapolated from adults using metabolic weight (body weight to the power of 0.75), which provided higher values than using isometric scaling (linear with body weight). This was to ensure a sufficient RDA, based on indications from studies conducted in developing countries that xerophthalmia and serum retinol concentrations of less than 200 $\mu\text{g/L}$ exist among preschool children with daily vitamin A intakes of up to 200 μg , whereas 300 $\mu\text{g/day}$ of vitamin A is associated with serum retinol concentrations greater than 300 $\mu\text{g/L}$ (Reddy, 1985). The RDA was set by using a CV of 20 %, as for adults.

For infants, Afssa considered an intake of 350 μg RE/day, based on a breast milk concentration of 500 μg RE/L and an ingested volume of 750 mL/day. For children, Afssa (2001) extrapolated the data from adults based on energy requirements.

The SCF (1993) proposed a PRI of 350 μg RE/day for infants aged 6–11 months based on the amount of vitamin A ingested with breast milk (FAO/WHO, 1988). PRIs for older children were set to make a smooth transition from infant to adult values. Although it was considered that there is little evidence to support these values, they appeared unlikely to be underestimates. A daily intake of about 300 μg has been reported to meet requirements of preschool children (Reddy, 1985).

For infants aged 6–11 months, the Netherlands Food and Nutrition Council (1992) set an AI on the basis of the concentration of vitamin A in breast milk. For children and adolescents, AIs were calculated by interpolation from the values for infants and adults, and an allowance was made for body weight and growth.

For infants, the UK COMA (DH, 1991) adopted the approach proposed by FAO/WHO (1988), which is described above. Values for children were interpolated from the values for infants and adults.

Table 2: Overview of Dietary Reference Values for vitamin A for infants and children

	NCM (2014)	D-A-CH (2013)	WHO/FAO (2004)	Afssa (2001)	IOM (2001)	SCF (1993)	NL (1992)	DH (1991)
Age (months)	6–11	4–12	7–12	0–12	7–12	6–11	6–11	7–12
PRI (µg RE/day)	300	600	400 ^(a)	350	500 ^(b)	350	400 ^(c)	350
Age (years)	1–2	1–4	1–3	1–3	1–3	1–3	1–4	1–3
PRI (µg RE/day)	300	600	400 ^(a)	400	300 ^(b)	400	400 ^(c)	400
Age (years)	2–5	4–7	4–6	4–6	4–8	4–6	4–7	4–6
PRI (µg RE/day)	350	700	450 ^(a)	450	400 ^(b)	400	500 ^(c)	400
Age (years)	6–9	7–10	7–9	7–9	9–13	7–10	7–10	7–10
PRI (µg RE/day)	400	800	500 ^(a)	500	600 ^(b)	500	700 ^(c)	500
Age (years)	10–13	10–13	10–18	10–12	14–18	11–14	10–13	11–14
PRI Boys (µg RE/day)	600	900	600 ^(a)	550	900 ^(b)	600	1 000 ^(c)	600
PRI Girls (µg RE/day)	600	900	600 ^(a)	550	700 ^(b)	600	800 ^(c)	600
Age (years)	14–17	13–15		13–15		15–17	13–16	15–18
PRI Boys (µg RE/day)	700	1 100		700		700	1 000 ^(c)	700
PRI Girls (µg RE/day)	700	1 000		600		600	800 ^(c)	600
Age (years)		15–19		16–19			16–19	
PRI Boys (µg RE/day)		1 100		800			1 000 ^(c)	
PRI Girls (µg RE/day)		900		600			800 ^(c)	

NCM, Nordic Council of Ministers; NL, Netherlands Foods and Nutrition Council; PRI, Population Reference Intake; RE, Retinol Equivalent; RAE, Retinol Activity Equivalent.

(a): Recommended Safe Intake.

(b): Expressed in µg RAE/day.

(c): Adequate Intake.

4.3. Pregnancy

The Nordic countries considered a retinol accumulation of 50 µg/day in the fetus and set a recommended intake of 800 µg RE/day for pregnant women to cover individual variation (Nordic Council of Ministers, 2014).

D-A-CH (2013) estimated that pregnant women should ingest on average one-third more than non-pregnant women, and a recommended intake of 1 100 µg RE/day was proposed throughout pregnancy.

WHO/FAO (2004) considered that a newborn infant appears to require around 100 µg retinol/day to meet its needs for normal growth and assumed that the fetus has similar needs during the third trimester of pregnancy. Recognising that a large portion of the world's population of pregnant women live under conditions of deprivation, an increment of 200 µg RE/day was proposed during the whole period of pregnancy, in order to enhance maternal storage during early pregnancy and to cover the needs of the rapidly growing fetus in late pregnancy. Finally, the Recommended Safe Intake was increased by 300 µg RE/day compared to that of non-pregnant women aged 19–65 years.

The IOM used a model based on the accumulation of vitamin A in the liver of the fetus during gestation and assuming that the liver contains approximately half of the body's amount of vitamin A when liver stores are low, as is the case for newborns (IOM, 2001). A concentration of 3 600 µg per fetus was calculated. Assuming the efficiency of maternal vitamin A absorption to average 70 % and vitamin A to be accumulated mostly in the last 90 days of pregnancy, the maternal requirement would be increased by around 50 µg/day during the last trimester. As vitamin A in the maternal diet may be stored and mobilised later as needed and some vitamin A may be retained in the placenta, the IOM proposed an additional requirement of 50 µg RAE/day for the entire pregnancy. The RDA was set by using a CV of 20 %, as for non-pregnant adults.

Afssa (2001) noted that fetal requirements are low and low amounts of vitamin A are accumulated in fetal liver. An increase of the recommended intake to 700 µg RE/day during the last trimester of pregnancy was proposed.

The SCF (1993) proposed a PRI of 700 µg RE/day during pregnancy, in order to enhance maternal storage to provide adequate vitamin A for the growing fetus in late pregnancy.

The Netherlands Food and Nutrition Council (1992) proposed an additional intake of 200 µg RE/day during pregnancy, based on fetal needs during the last three months of pregnancy (Olson and Hodges, 1987).

The UK COMA considered that an increment of 100 µg RE/day during pregnancy should enhance maternal storage and allow adequate vitamin A for the growing fetus in late pregnancy (DH, 1991).

Table 3: Overview of Dietary Reference Values for vitamin A for pregnant women

	NCM (2014)	D-A-CH (2013)	WHO/FAO (2004)	Afssa (2001)	IOM (2001)	SCF (1993)	NL (1992)	DH (1991)
Age (years)					14–18			
PRI (µg RE/day)	800	1 100	800 ^(a)	700 ^(b)	750 ^(c)	700	1 000 ^(d)	700
Age (years)					≥ 19			
PRI (µg RE/day)					770 ^(c)			

NCM, Nordic Council of Ministers; NL, Netherlands Foods and Nutrition Council; PRI, Population Reference Intake; RE, Retinol Equivalent; RAE, Retinol Activity Equivalent.

(a): Recommended Safe Intake.

(b): Third trimester of pregnancy.

(c): Expressed in µg RAE/day.

(d): Adequate Intake.

4.4. Lactation

The Nordic countries proposed an additional intake of 400 µg RE/day for lactating women, to compensate for the amount of vitamin A secreted in breast milk and considering reported values for breast milk vitamin A concentration of 450–600 µg RE/L in Western countries and an average milk production of 750 mL/day (Nordic Council of Ministers, 2014).

D-A-CH (2013) noted that the intake of breast-fed infants is about 500 µg RE/day (Souci et al., 2000). With prolonged breastfeeding, the vitamin A concentration of breast milk decreases while the breast-fed infant requires additional vitamin A for growth. Mainly for women breastfeeding longer than four months an allowance of 700 µg RE/day was recommended to satisfy the infant's requirement and to avoid deficits in the mother.

WHO/FAO proposed an increment of 350 µg RE/day to replace the amount secreted in breast milk (WHO/FAO, 2004).

The IOM considered that a breast-fed infant ingests on average 400 µg RAE/day in the first six months of life, and this was proposed as the additional EAR during lactation to maintain adequate vitamin A body stores of the mother (IOM, 2001). The RDA was set by using a CV of 20 %.

Afssa considered that breastfeeding women secrete around 350 µg RE/day (based on a concentration of 500 µg RE/L and a secreted amount of 750 mL/day) and this was proposed as the additional amount during lactation to be added to the PRI for non-lactating women (Afssa, 2001).

The SCF assumed that 350 µg RE/day is supplied in breast milk and proposed an increment of this amount throughout lactation (SCF, 1993).

The Netherlands Food and Nutrition Council (1992) recommended an additional intake of 450 µg RE/day during lactation, to offset the loss of vitamin A through breast milk, assuming an average concentration of 550 µg/L.

The UK COMA proposed an increment of 350 µg RE/day during lactation to cover the amount of vitamin A secreted with breast milk (DH, 1991).

Table 4: Overview of Dietary Reference Values for vitamin A for lactating women

	NCM (2014)	D-A-CH (2013)	WHO/FAO (2004)	Afssa (2001)	IOM (2001)	SCF (1993)	NL (1992)	DH (1991)
Age (years)					14–18			
PRI (µg RE/day)	1 100	1 500	850 ^(a)	950	1 200 ^(b)	950	1 250 ^(c)	950
Age (years)					≥ 19			
PRI (µg RE/day)					1 300 ^(b)			

NCM, Nordic Council of Ministers; NL, Netherlands Foods and Nutrition Council; PRI, Population Reference Intake; RE, Retinol Equivalent; RAE, Retinol Activity Equivalent.

(a): Recommended Safe Intake

(b): Expressed in µg RAE/day.

(c): Adequate Intake

5. Criteria (endpoints) on which to base Dietary Reference Values

The average requirement of vitamin A is defined as the intake required to permit adequate growth and other vitamin A-dependent functions and to maintain an acceptable total body pool of the vitamin.

5.1. Indicators of vitamin A requirement

The requirement of vitamin A has been estimated by other expert bodies on the basis of the amount needed to correct deficiency symptoms such as impaired dark adaptation among vitamin A-depleted subjects (Netherlands Food and Nutrition Council, 1992; Afssa, 2001); to raise plasma retinol concentration into the normal range in depleted subjects (Netherlands Food and Nutrition Council, 1992; Afssa, 2001); and to maintain a given body pool size of retinol in well-nourished subjects (SCF, 1993; IOM, 2001; WHO/FAO, 2004; Nordic Council of Ministers, 2014).

5.1.1. Symptoms of vitamin A deficiency

Xerophthalmia is the most specific clinical consequence of vitamin A deficiency (Section 2.2.2.1). Markers of visual function have been developed to assess vitamin A status (Section 2.4.3). However, data relating such measurements to dietary vitamin A intake are scarce. In a depletion–repletion study in eight men in whom vitamin A deficiency was induced, daily supplementation with around 300 µg retinol (4–5 µg/kg body weight) corrected abnormalities in adaptation to dark and electroretinographic patterns (Saubberlich et al., 1974). This amount may be considered the minimal dietary requirement in adults to maintain normal visual function. However, it is recognised that the prevalence of ocular manifestations (i.e. xerophthalmia) often underestimates the magnitude of functional vitamin A deficiency. Therefore, such amount may not cover other vitamin A-dependent functions (Section 2.2.1) and allow an adequate total body pool of the vitamin to be maintained.

The Panel considers that these indicators cannot be used for deriving DRVs for vitamin A.

5.1.2. Plasma/serum retinol concentration

Plasma/serum retinol concentration lacks sensitivity and specificity as a marker of vitamin A status in the general healthy population, because of the tight homeostatic control of retinol concentration over the range of adequate liver retinol concentrations and the influence of a number of confounding factors (Section 2.4.2).

The Panel considers that this marker cannot be used for deriving DRVs for vitamin A.

5.1.3. Maintenance of total body retinol content and liver retinol concentration

Liver retinol concentration is a biomarker of vitamin A status. A concentration of 20 µg retinol/g liver (0.07 µmol/g) in adults represents a level assumed to maintain adequate plasma retinol concentrations, to prevent clinical signs of deficiency and to provide adequate stores (Section 2.4.1). Accordingly, the Panel considers that a concentration of 20 µg retinol/g liver (0.07 µmol/g) can be used as a target value for establishing the requirement for vitamin A in adults. In the absence of specific data for infants and children, the Panel considers that the same target value as for adults can be used in these age groups.

The dietary intake of vitamin A required to maintain this liver concentration can be determined from a factorial approach (Olson, 1987). Data on the relationship between dietary intake of vitamin A and retinol liver (or total body) stores measured by stable isotope dilution methods may also be used (Haskell et al., 2005) (Section 2.4.1.2).

5.1.3.1. Factorial approach

The vitamin A intake required to maintain a concentration of 20 µg retinol/g liver (0.07 µmol/g) can be calculated on the basis of the factorial approach proposed by Olson (1987), which takes into account the ratio of total body/liver retinol stores, the fractional catabolic rate of retinol and the efficiency of storage of ingested retinol.

To apply the factorial approach, a number of assumptions have to be made:

- The body store of retinol appears to be an important determinant of the catabolic rate of retinol (Section 2.3.6.1). Limited data are available on the fractional catabolic rate in subjects with adequate retinol body stores. Recent data indicate that the fractional catabolic rate may be higher than the value of 0.5 % which has usually been considered. Taking a conservative approach, the Panel assumes a fractional catabolic rate of 0.7 % for adults using the highest value of the range measured in four US adults at steady state (Section 2.3.6.1).
- In healthy individuals with an adequate vitamin A status, 70–90 % of body retinol is stored in the liver (Section 2.3.4.1). The Panel notes the paucity of data in humans. The Panel assumes a ratio of 80 % for all age groups.
- Available data indicate that the average efficiency of storage of retinol in the liver of adult subjects with adequate hepatic stores (≥ 20 µg retinol/g liver) is 42 %. Assuming that liver stores represent 80 % of the whole body stores in this population group, this would correspond to a storage efficiency in the whole body of 52 % (Section 2.3.4.3). The Panel assumes an efficiency of storage of retinol in the whole body of 50 % for all age groups.
- Based on available data which show that the liver/body weight ratio decreases with age (Haddad et al., 2001; Young et al., 2009), the Panel assumes average liver/body weight ratios of 4.0 % up to 3 years, 3.5 % from 4 to 6 years, 2.8 % from 7 to 14 years and 2.4 % above 15 years of age and in adults.
- The Panel considers that maintenance needs for vitamin A expressed with respect to body weight are the same for adults and children. A growth component has to be added for children to take into account higher vitamin A utilisation for growth needs (Section 2.3.6.1). Growth factors were calculated as the proportional increase in protein requirement for growth relative to the maintenance requirement at the different ages, as follows: 0.57 for infants aged 7–11 months, 0.25 for boys and girls aged 1–3 years, 0.06 for boys and girls aged 4–6 years, 0.13 for boys and girls aged 7–10 years, 0.11 for boys and 0.08 for girls aged 11–14 years, and 0.08 for boys and 0.03 for girls aged 15–17 years (EFSA NDA Panel, 2014a).

The Panel notes that data on total body/liver retinol stores in humans are scarce, that available information on the fractional catabolic rate and efficiency of storage of retinol comes from studies involving a small number of subjects and that the influence of factors such as age is not well characterised.

5.1.3.2. Data from stable isotope dilution methods

Haskell et al. (2011) investigated the amount of daily vitamin A required to maintain liver stores in a selected population of Bangladeshi men expected to have a liver retinol concentration close to 20 µg/g (0.07 µmol/g). During a 60-day intervention period, 16 subjects (18–32 years, body weight ~50 kg) consumed a controlled basal diet containing 100 µg RAE/day and were randomly assigned to receive one of eight different amounts of retinol (range 100–1 000 µg/day; n = 2 per group) in the form of retinyl palmitate dissolved in corn oil. The total body retinol pool sizes and liver retinol concentrations were quantitatively estimated by using the DRD method before and after the intervention. A “semi-quantitative” estimate of the change in retinol pool size was also obtained by estimating the change in plasma isotopic ratios on day 3 after dosing, before and after the intervention. Mean (± SD) estimated total body retinol pool sizes were 17 ± 9 mg (59 ± 32 µmol) at baseline and 18 ± 10 mg (64 ± 34 µmol) after the intervention, and liver retinol concentrations were 13 ± 7 µg/g (0.047 ± 0.025 µmol/g) and 14 ± 8 µg/g (0.049 ± 0.027 µmol/g), respectively. There were significant linear relationships between daily supplemental retinol intake and the changes in total body retinol pool size as assessed quantitatively (r = 0.62, p = 0.010) or “semi-quantitatively” (r = 0.68, p = 0.004). From the respective regression lines, the authors estimated that a daily supplement of 400 µg retinol (95 % CI = undefined–640) with the quantitative approach, and of 254 µg/day (95 % CI = 156–336) with the “semi-quantitative” approach, would be required to maintain the retinol pool size of 17 mg (60 µmol) (13 ± 7 µg/g liver (0.047 ± 0.025 µmol/g liver)). Considering the background dietary intake, vitamin A intakes of 500 or 354 µg RAE/day were derived from the two methods. The Panel notes that the estimated liver retinol concentration in the study population was lower than the target of 20 µg/g liver (0.07 µmol/g liver). The authors indicate that no signs or symptoms of vitamin A deficiency were identified in the subjects, but the publication does not provide details on the physical examination undertaken, including eye/vision assessment.

Ribaya-Mercado et al. (2004) investigated the relationship between dietary vitamin A intake and total body and liver retinol stores in a cross-sectional study in men (n = 31, body weight 53.3 ± 9.7 kg) and women (n = 31, body weight 45.9 ± 10.1 kg) aged 60–88 years in the rural Philippines. Total body pool was assessed using the DRD method and vitamin A intake was estimated by three non-consecutive 24-hour dietary recalls. Mean (± SD) (range) estimated retinol pool size was 75 ± 41 mg (11–190 mg) (263 ± 144 µmol (38–664 µmol)) in men and 62 ± 39 mg (6–169 mg) (215 ± 137 µmol (20–590 µmol)) in women. Assuming that liver weight was 2.4 % of body weight in adults and that, in these marginally nourished individuals, 70 % of total body retinol was found in the liver, the authors estimated a mean (± SD) liver retinol concentration of 40 ± 17 (range 7–74) µg/g (0.139 ± 0.058 (range 0.026–0.260) µmol/g) in men and 40 ± 27 (range 5–125) µg/g (0.140 ± 0.095 (range 0.019–0.438) µmol/g) in women. The mean vitamin intake of the men and women with a liver concentration ≥ 20 µg retinol/g (0.07 µmol/g) (n = 53) was 135 ± 86 µg RAE/day (n = 27) and 134 ± 104 µg RAE/day (n = 26), respectively.

Valentine et al. (2013) assessed the relationship between vitamin A intake and total body retinol pool size in another cross-sectional study in 40 non-pregnant, non-lactating US women (22.4 ± 2.3 years, body weight 61.2 ± 7.2 kg). Body pool size and liver stores were estimated by using a [¹³C₂]-RID test. Mean (± SD) estimated total body retinol pool size was 234 ± 154 mg (range 41–893 mg) (816.5 ± 537.4 µmol (range 141.5–3 116 µmol)). Eighty per cent of total body retinol was assumed to be found in the liver and the liver weight was assumed to represent 2.4 % of body weight. Estimated mean liver retinol concentration was 129 ± 89 µg/g (0.45 ± 0.31 µmol/g) and ranged from 26 µg/g (0.09 µmol/g) to 513 µg/g (1.79 µmol/g). Mean (± SD) vitamin A intake as assessed by food frequency questionnaire (FFQ) (including supplements) was 1 213 ± 778 µg RAE/day (range 378–3 890 µg RAE/day) and was positively correlated with liver retinol concentration and total body

retinol pool size (Pearson correlation coefficients 0.41 and 0.40, $p = 0.009$ and $p = 0.011$, respectively). Vitamin A intake was also estimated by a three-day dietary record; mean (\pm SD) estimate was $1\,180 \pm 705$ μg RAE/day (range 78–3 020 μg RAE/day) and no significant correlation was found with liver retinol concentration and total body retinol pool size. In a subset of women with a mean daily vitamin A intake (521 ± 119 μg RAE/day) similar to the EAR set by the IOM (2001) on the basis of the Olson equation and a target liver concentration of 20 μg retinol/g (0.07 $\mu\text{mol/g}$), the authors found an average liver retinol concentration of 86 ± 29 $\mu\text{g/g}$ (0.30 ± 0.10 $\mu\text{mol/g}$).

In a group of 32 young women (19–30 years) in the USA with a mean (\pm SD) vitamin A intake of $1\,148 \pm 782$ μg RAE (assessed by FFQ, including supplements), Valentine (2013) estimated a mean total body retinol pool size of 234 ± 158 mg (817 ± 550 μmol) using the [$^{13}\text{C}_2$]-RID test. A mean liver retinol concentration of 132 ± 92 $\mu\text{g/g}$ (0.46 ± 0.32 $\mu\text{mol/g}$) was derived. Participants consumed a study diet providing 175 μg (0.6 μmol) RAE/day for 12 weeks. For the middle six weeks (day 14 to day 56), women were randomised to take a daily supplement of 0, 175 μg or 525 μg retinol as retinyl palmitate. No significant changes in liver retinol concentration and total body retinol pool size between and within groups were found after the intervention.

The Panel notes that current data on the dose–response relationship between vitamin A intake and liver stores are limited and difficult to compare owing to differences in the vitamin A status of the study populations and in study designs. The Panel also notes uncertainties related to the quantitative total body pool and liver store estimates derived from the stable isotope dilution methods, owing to the different assumptions made, and uncertainties related to vitamin A intake estimates inherent to the dietary assessment methods used and the conversion of provitamin A carotenoids into RE. Despite these uncertainties, the Panel notes that two studies (Ribaya-Mercado et al., 2004; Valentine et al., 2013) suggest that the amount of dietary vitamin A required to achieve a target liver retinol concentration of 20 $\mu\text{g/g}$ (0.07 $\mu\text{mol/g}$) may be lower than previously calculated on the basis of the equation proposed by Olson (1987).

The Panel considers that the available data from stable isotope methods are insufficient to derive the requirement for vitamin A for adults.

5.2. Indicators of vitamin A requirement in pregnancy and lactation

During pregnancy there is an additional need of vitamin A for the fetus and possibly for the growth of maternal tissues. However, data are scarce.

Based on data from Thai fetuses of healthy mothers with an average retinol content in the liver of 1 800 μg (6 μmol) at 37–40 gestational weeks ($n = 10$) (Montreewasuwat and Olson, 1979) and assuming that the liver contains approximately half of the body's retinol content when liver stores are low, as is the case for newborns, a total amount of retinol of 3 600 μg (12 μmol) in the fetus was estimated by IOM (2001).

There is no information on the amount of retinol accumulated in maternal tissue formed during pregnancy.

With respect to lactating women, the Panel estimated a retinol secretion of 424 $\mu\text{g/day}$ in breast milk during the first six months of lactation (Section 2.3.6.3). As discussed by Allen and Haskell (2001), liver stores at birth are low and sufficient to supply the infant's vitamin A requirement for only a few days, even when the mother is well nourished during pregnancy. Allen and Haskell (2001) assumed that an intake of about 300 μg retinol/day is required to accumulate adequate liver stores during the first six months of life, although analytical data to support this are lacking.

The Panel considers that data on whole body retinol stores in the fetus and on retinol secretion in breast milk can be used to derive the additional requirement for, respectively, pregnant or lactating women.

5.3. Vitamin A intake and health consequences

A comprehensive search of the literature published between 1 January 1990 and 1 July 2011 was performed as preparatory work to identify relevant health outcomes on which DRVs for vitamin A may be based (Heinonen et al., 2012). Additional literature searches were performed until October 2014.

A number of intervention studies in children have assessed the effect of vitamin A supplementation on the risk of (premature) death, and the incidence and severity of diarrhoea, measles and lower respiratory tract infections (Fawzi et al., 1992; Anonymous, 1993; Beaton et al., 1993; Glasziou and Mackerras, 1993; Grotto et al., 2003; Brown and Roberts, 2004; Wu et al., 2005; Chen et al., 2008; Imdad et al., 2011; Mayo-Wilson et al., 2011; McLaren and Kraemer, 2012). In adults, intervention studies have investigated the effect of supplementation with retinol, often in combination with other nutrients, for the primary prevention of a variety of diseases, including cancer of various sites (Bjelakovic et al., 2006; Bjelakovic et al., 2008; Misotti and Gnagnarella, 2013) and reproduction-related outcomes (Thorne-Lyman and Fawzi, 2012), and in relation to all-cause mortality (Fortmann et al., 2013; Bjelakovic et al., 2014). The Panel notes that these studies typically used high doses of vitamin A (1 000–60 000 µg RE in daily or bolus doses) and that background vitamin A intake was not assessed in these studies. The Panel considers that these intervention studies cannot be used for the setting of DRVs for vitamin A.

The relationship between vitamin A intake and health outcomes has been investigated in observational (case-control, cross-sectional, prospective cohort) studies, in which an association between vitamin A intake and a health outcome might be confounded by uncertainties inherent in the methodology used for the assessment of vitamin A intake, and by the effect of other dietary, lifestyle, or undefined factors on the disease outcomes investigated. The Panel notes that different definitions of “vitamin A” have been applied among studies (i.e. defined as retinol only or as retinol and provitamin A carotenoids expressed in IU, µg RE or µg RAE, or undefined).

No association was observed between retinol intake and all-cause or cardiovascular disease mortality in a cohort study in the UK (Fletcher et al., 2003), or between intake of “vitamin A” or retinol and risk of death from coronary heart disease in the prospective Iowa Women’s Health study (Kushi et al., 1996).

Several studies reported on the association between intake of “vitamin A” or retinol and risk of cancer at various sites, including risk of oral premalignant lesions (one prospective cohort study (Maserejian et al., 2007)), nasopharyngeal carcinoma (one case-control study (Hsu et al., 2012)), lung cancer (two prospective cohort studies (Yong et al., 1997; Takata et al., 2013)), benign proliferative epithelial disorders of the breast (one case-control study (Rohan et al., 1990); one nested case-control study (Rohan et al., 1998)), breast cancer (one meta-analysis (Fulan et al., 2011)), gastric cancer (two prospective cohort studies (Larsson et al., 2007; Miyazaki et al., 2012)), pancreatic cancer (two case-control studies (Zablotska et al., 2011; Jansen et al., 2013)), colorectal cancer (three case-control studies (Key et al., 2012; Wang et al., 2012; Leenders et al., 2014); one prospective cohort study (Ruder et al., 2011); one systematic review (Xu et al., 2013)), prostate cancer (one case-control study (Ghadirian et al., 1996); one prospective cohort study (Giovannucci et al., 1995)), cervical cancer (two systematic reviews (Garcia-Closas et al., 2005; Zhang et al., 2012); one prospective cohort study (Gonzalez et al., 2011)), ovarian cancer (one case-control study (Zhang et al., 2004); one prospective cohort study (Fairfield et al., 2001)), bladder cancer (one case-control study (Garcia-Closas et al., 2007)), melanoma or basal cell carcinoma (one case-control study (Naldi et al., 2004); three prospective cohort studies (Fung et al., 2002; Feskanich et al., 2003; Asgari et al., 2012)) and non-Hodgkin’s lymphoma (one case-control study (Mikhak et al., 2012); one prospective cohort study (Kabat et al., 2012)). Results were limited and/or inconsistent.

Some observational studies have assessed the association between “vitamin A” or retinol intake and asthma, wheeze or other measures of lung function with inconsistent results (a systematic review

including two prospective cohort studies, one nested case–control study, ten case–control studies and six cross-sectional studies (Allen et al., 2009); a systematic review including two case–control and three cross-sectional studies (Nurmatov et al., 2011); a prospective cohort study (Maslova et al., 2014)).

Some observational studies investigated the association between “vitamin A” or retinol intake and eye health-related outcomes, including cataract (one cross-sectional study (Cumming et al., 2000); one prospective study (Chasan-Taber et al., 1999)), age-related maculopathy (one cross-sectional study (Smith et al., 1999)) and age-related macular degeneration (one case–control study (Seddon et al., 1994)) and glaucoma (one cross-sectional study (Giacconi et al., 2012); two cohort studies (Kang et al., 2003; Ramdas et al., 2012)). Results were limited and/or inconsistent.

In view of the limited and/or inconsistent evidence on an association between vitamin A or retinol intake and these health outcomes, the Panel considers that the data available cannot be used for setting DRVs for vitamin A.

6. Data on which to base Dietary Reference Values

The Panel expresses DRVs for vitamin A in $\mu\text{g RE/day}$ (Section 2.3.8). The requirement of vitamin A can be met with any mixture of preformed vitamin A and provitamin A carotenoids that provides an amount of vitamin A equivalent to the reference value in terms of $\mu\text{g RE/day}$.

6.1. Adults

The Panel determines the AR for vitamin A in healthy adults as the vitamin A intake required to maintain a liver retinol concentration of $20 \mu\text{g/g}$ ($0.07 \mu\text{mol/g}$). The latter is considered by the Panel as indicative of an adequate vitamin A status (or vitamin A body pool) at which the different functions of vitamin A in the body can be fulfilled (Sections 2.4.1, 2.4.4 and 5.1.3).

In the absence of a better characterisation of the relationship between dietary intake of vitamin A and liver stores, the requirement to maintain a concentration of $20 \mu\text{g retinol/g liver}$ ($0.07 \mu\text{mol/g}$) can be calculated on the basis of the factorial approach as proposed by Olson (1987), as follows:

$$\text{AR } (\mu\text{g RE/day}) = \text{target liver concentration } (\mu\text{g retinol/g}) \times \text{body/liver retinol stores ratio} \times \text{liver/body weight ratio } (\%) \times \text{fractional catabolic rate of retinol } (\%) \times (1/\text{efficiency of body storage } (\%)) \times \text{reference body weight } (\text{kg}) \times 10^3$$

The Panel uses the following values for adults (Section 5.1.3.1): (1) a total body/liver retinol store ratio of 1.25 (i.e. 80 % of vitamin A in the body is stored in the liver); (2) a liver/body weight ratio of 2.4 %; (3) a fractional catabolic rate of retinol of 0.7 % per day; and (4) an efficiency of storage in the whole body of ingested retinol of 50 %. Reference body weights for adult women and men in the EU are 58.5 and 68.1 kg, respectively (EFSA NDA Panel, 2013).

On the basis of this calculation, ARs of $570 \mu\text{g RE/day}$ for men and $490 \mu\text{g RE/day}$ for women are derived after rounding.

Assuming a CV of 15 % because of the variability in requirement and of the large uncertainties in the dataset (see Section 5.1.3.1), PRIs of $750 \mu\text{g RE/day}$ for men and $650 \mu\text{g RE/day}$ for women are set. PRIs were rounded to the closest 50 or 100.

Table 5: Dietary Reference Values for vitamin A for men and women

Reference body weight ^(a) (kg)		AR (µg RE/day) ^(b)		PRI (µg RE/day) ^(c)	
Men	Women	Men	Women	Men	Women
68.1	58.5	570	490	750	650

(a): Median body weight of 18- to 79-year-old men and women, respectively, based on measured body heights of 16 500 men and 19 969 women in 13 EU Member States and assuming a body mass index (BMI) of 22 kg/m²; see Appendix 11 in EFSA NDA Panel (2013)).

(b): Values for ARs were rounded to the closest 10.

(c): Values for PRIs were rounded to the closest 50, but PRIs were calculated based on the unrounded ARs.

6.2. Infants and children

Breast milk retinol concentration is influenced by maternal vitamin A status, and large variation in mean total retinol concentration of mature breast milk is observed (Section 2.3.6.3). Using estimated retinol intakes in exclusively breast-fed infants as a basis for setting a DRV for infants aged 7–11 months is therefore associated with considerable uncertainty. Thus, the Panel considers it more appropriate to derive DRVs for infants aged 7–11 months on the same basis as for adults.

For infants aged 7–11 months and children, the average intake of vitamin A required to maintain a concentration of 20 µg retinol/g liver is calculated with the same equation as for adults but with values for reference body weight and liver/body weight ratio specific to infants and children (Section 5.1.3.1). Although there is some indication that the fractional catabolic rate of retinol may be higher in children than in adults, data are limited (Section 2.3.6.1). In the absence of more robust data, the Panel decides to apply the value for fractional catabolic rate observed in adults and to correct it on the basis of a growth factor (Section 5.1.3.1).

This approach is preferred to scaling down from adults based on body weight (either isometric or allometric), as retinol is mainly stored in the liver, the size of which does not linearly change with body weight during growth, and as vitamin A requirement is not directly related to energy needs and expenditure.

The requirement to maintain a concentration of 20 µg retinol/g liver can be calculated in infants and children on the basis of the factorial approach as follows:

$$\text{AR } (\mu\text{g RE/day}) = \text{target liver concentration } (\mu\text{g retinol/g}) \times \text{body/liver retinol stores ratio} \times \text{liver/body weight ratio } (\%) \times \text{fractional catabolic rate of retinol } (\%) \times (1/\text{efficiency of body storage } (\%)) \times \text{reference body weight } (\text{kg}) \times (1 + \text{growth factor}) \times 10^3$$

The Panel uses the following values for infants aged 7–11 months and children (Section 5.1.3.1): (1) a total body/liver retinol stores ratio of 1.25 (i.e. 80 % of retinol in the body is stored in the liver); (2) an age-specific liver/body weight ratio; (3) a fractional catabolic rate of retinol of 0.7 % per day; (4) an efficiency of storage in the whole body of ingested retinol of 50 %; and (5) a growth factor of 0.57 for infants aged 7–11 months, 0.25 for boys and girls aged 1–3 years, 0.06 for boys and girls aged 4–6 years, 0.13 for boys and girls aged 7–10 years, 0.11 for boys and 0.08 for girls aged 11–14 years, and 0.08 for boys and 0.03 for girls aged 15–17 years (EFSA NDA Panel, 2014a).

It was considered unnecessary to give sex-specific values for infants and children up to 14 years of age. As for adults, a CV of 15 % is used for setting PRIs for the respective age groups (Table 6). PRIs were rounded to the closest 50 or 100.

Table 6: Dietary Reference Values for vitamin A for infants and children

Age	Reference body weight (kg)	Liver weight (% body weight)	Growth factor	AR ^(h) (µg RE/day)	PRI ⁽ⁱ⁾ (µg RE/day)
7–11 months	8.6 ^(a)	4.0	0.57	190	250
1–3 years	11.9 ^(b)	4.0	0.25	205	250
4–6 years	19.0 ^(c)	3.5	0.06	245	300
7–10 years	28.7 ^(d)	2.8	0.13	320	400
11–14 years	44.6 ^(e)	2.8	0.11 (M) / 0.08 (F)	480	600
15–17 years (M)	64.1 ^(f)	2.4	0.08	580	750
15–17 years (F)	56.4 ^(g)	2.4	0.03	490	650

AR, Average Requirement; PRI, Population Reference Intake; F, females; M, males.

(a): Mean of the body weight-for-age at 50th percentile of male or female infants aged nine months according to the WHO Growth Standards (WHO Multicentre Growth Reference Study Group, 2006).

(b): Mean of body weight-for-age at 50th percentile of boys and girls aged 24 months (WHO Multicentre Growth Reference Study Group, 2006).

(c): Mean of body weight at 50th percentile of boys and girls aged 5 years (van Buuren et al., 2012).

(d): Mean of body weight at 50th percentile of boys and girls aged 8.5 years (van Buuren et al., 2012).

(e): Mean of body weight at 50th percentile of boys and girls aged 12.5 years (van Buuren et al., 2012).

(f): Body weight at 50th percentile of boys aged 16 years (van Buuren et al., 2012).

(g): Body weight at 50th percentile of girls aged 16 years (van Buuren et al., 2012).

(h): Values for ARs were rounded to the closest 5 or 10.

(i): Values for PRIs were rounded to the closest 50 or 100, but PRIs were calculated based on the unrounded ARs.

6.3. Pregnancy

The Panel assumes that a total amount of 3 600 µg retinol is accumulated in the fetus over the course of pregnancy (Section 5.2). Considering that the accretion mostly occurs in the last months of pregnancy, and assuming an efficiency of storage of 50 % for the fetus, an additional daily requirement of 51 µg RE is calculated for the second half of pregnancy (i.e. 3 600 µg/140 days × 2). In order to allow for the extra need related to the growth of maternal tissues (e.g. placenta), the Panel applies this additional requirement to the whole period of pregnancy.

Consequently, an AR of 540 µg RE/day is estimated for pregnant women by adding the additional requirement of pregnancy to the AR for non-pregnant non-lactating women and rounding. Considering a CV of 15 % and rounding to the closest 100, a PRI of 700 µg RE/day is derived for pregnant women.

6.4. Lactation

Based on an average amount of retinol secreted in breast milk of 424 µg/day (Section 2.3.6.3) and an absorption efficiency of retinol of 80 % (Section 2.3.1.1), an additional vitamin A intake of 530 µg RE/day is considered sufficient to replace these losses. An AR of 1 020 µg RE/day is estimated by adding the additional requirement of lactation to the AR for non-pregnant non-lactating women. Considering a CV of 15 % and rounding to the closest 100, a PRI of 1 300 µg RE/day is proposed for lactating women.

CONCLUSIONS

The Panel concluded that ARs and PRIs for vitamin A in healthy adults can be derived from the vitamin A intake required to maintain a concentration of 20 µg retinol/g liver (0.07 µmol/g). In the absence of a better characterisation of the relationship between dietary intake of vitamin A and liver stores, ARs for adults were calculated on the basis of a factorial approach which takes into account the ratio of total body/liver retinol stores, the fractional catabolic rate of retinol and the efficiency of storage of ingested retinol. For infants aged 7–11 months and children, ARs were derived on the basis of the same equation as for adults, by using specific age-related values for reference body weight and liver/body weight ratio. For catabolic rate, the value for adults corrected on the basis of a growth factor was used. It was considered unnecessary to give sex-specific values for infants and children up to 14 years of age. The estimated amount of retinol accumulated in the fetus over the course of pregnancy was used as a basis to increase the AR for pregnant women. For lactating women, an

increase in the AR was based on the vitamin A intake required to compensate for the amount of retinol secreted in breast milk. Owing to the variability in requirement and the large uncertainties in the dataset, a CV of 15 % was used to calculate PRIs for all population groups (Table 7).

Table 7: Summary of Population Reference Intakes for vitamin A

Age	Population Reference Intake (µg/day)
7–11 months	250
1–3 years	250
4–6 years	300
7–10 years	400
11–14 years	600
15–17 years (M)	750
15–17 years (F)	650
≥ 18 years (M)	750
≥ 18 years (F)	650
Pregnancy	700
Lactation	1 300

F, females; M, males.

RECOMMENDATIONS FOR RESEARCH

The Panel recommends the following actions:

- pursue the characterisation of provitamin A carotenoid bioconversion to retinol;
- pursue the development of indirect measurement of liver stores by stable isotope dilution methods and application of the method to inform the dose–response relationship between vitamin A intake and retinol liver stores;
- further investigate and characterise the fractional catabolic rate of retinol and its determinants, including the influence of hepatic retinol stores, age and physiological state (e.g. pregnancy);
- characterise the efficiency of storage of a physiological dose of retinol in populations with adequate status;
- undertake research regarding the vitamin A requirement of infants and children;
- further characterise the relationship between vitamin A intake and health effects across the range of dietary intakes;
- further investigate the genetic basis of the differences in efficiency in provitamin A carotenoid and retinol metabolism in humans.

REFERENCES

- Abedin Z, Hussain MA and Ahmad K, 1976. Liver reserve of vitamin A from medico-legal cases in Bangladesh. *Bangladesh Medical Research Council Bulletin*, 2, 43-51.
- Abumrad N, Harmon C and Ibrahimi A, 1998. Membrane transport of long-chain fatty acids: evidence for a facilitated process. *Journal of Lipid Research*, 39, 2309-2318.
- Afssa (Agence française de sécurité sanitaire des aliments), 2001. Apports nutritionnels conseillés pour la population française. Editions Tec&Doc, Paris, France, 605 pp.
- Afssa (Agence française de sécurité sanitaire des aliments), 2009. Étude Individuelle Nationale des Consommations Alimentaires 2 (INCA 2) 2006-2007 (version 2). Maisons-Alfort, France, 225 pp.
- Al-Delaimy WK, van Kappel AL, Ferrari P, Slimani N, Steghens JP, Bingham S, Johansson I, Wallstrom P, Overvad K, Tjonneland A, Key TJ, Welch AA, Bueno-de-Mesquita HB, Peeters PH, Boeing H, Linseisen J, Clavel-Chapelon F, Guibout C, Navarro C, Quiros JR, Palli D, Celentano E, Trichopoulou A, Benetou V, Kaaks R and Riboli E, 2004. Plasma levels of six carotenoids in nine European countries: report from the European Prospective Investigation into Cancer and Nutrition (EPIC). *Public Health Nutrition*, 7, 713-722.
- Al-Mekhlafi HM, Al-Zabedi EM, Al-Maktari MT, Atroosh WM, Al-Delaimy AK, Moktar N, Sallam AA, Abdullah WA, Jani R and Surin J, 2013. Effects of vitamin A supplementation on iron status indices and iron deficiency anaemia: a randomized controlled trial. *Nutrients*, 6, 190-206.
- Al Tanoury Z, Piskunov A and Rochette-Egly C, 2013. Vitamin A and retinoid signaling: genomic and nongenomic effects. *Journal of Lipid Research*, 54, 1761-1775.
- Allen LH and Haskell M, 2001. Vitamin A requirements of infants under six months of age. *Food and Nutrition Bulletin*, 22, 214-234.
- Allen S, Britton JR and Leonardi-Bee JA, 2009. Association between antioxidant vitamins and asthma outcome measures: systematic review and meta-analysis. *Thorax*, 64, 610-619.
- Altmann SW, Davis HR, Jr., Zhu LJ, Yao X, Hoos LM, Tetzloff G, Iyer SP, Maguire M, Golovko A, Zeng M, Wang L, Murgolo N and Graziano MP, 2004. Niemann-Pick C1 Like 1 protein is critical for intestinal cholesterol absorption. *Science*, 303, 1201-1204.
- Ambrosini GL, Bremner AP, Reid A, Mackerras D, Alfonso H, Olsen NJ, Musk AW and de Klerk NH, 2013. No dose-dependent increase in fracture risk after long-term exposure to high doses of retinol or beta-carotene. *Osteoporosis International*, 24, 1285-1293.
- Ambrosini GL, Alfonso H, Reid A, Mackerras D, Bremner AP, Beilby J, Olsen NJ, Musk AW and de Klerk NH, 2014. Plasma retinol and total carotenes and fracture risk after long-term supplementation with high doses of retinol. *Nutrition*, 30, 551-556.
- Amcoff E, Edberg A, Enghardt Barbieri H, Lindroos A-K, Nälsén C, Pearson M and Warensjö Lemming E, 2012. [Riksmaten – vuxna 2010–11. Livsmedels- och näringsintag bland vuxna i Sverige. Resultat från matvaneundersökning utförd 2010–11]. Livsmedelsverket, Uppsala, Sverige, 180 pp.
- Amedee-Manesme O, Mourey MS, Hanck A and Therasse J, 1987. Vitamin A relative dose response test: validation by intravenous injection in children with liver disease. *American Journal of Clinical Nutrition*, 46, 286-289.
- Amedee-Manesme O, Luzeau R, Wittepen JR, Hanck A and Sommer A, 1988. Impression cytology detects subclinical vitamin A deficiency. *American Journal of Clinical Nutrition*, 47, 875-878.
- Amine EK, Corey J, Hegsted DM and Hayes KC, 1970. Comparative hematology during deficiencies of iron and vitamin A in the rat. *Journal of Nutrition*, 100, 1033-1040.

- Anonymous, 1983. Nomenclature of retinoids: recommendations 1981. IUPAC-IUB Joint Commission on Biochemical Nomenclature (JCBN). Archives of Biochemistry and Biophysics, 224, 728-731.
- Anonymous, 1993. Vitamin A supplementation in northern Ghana: effects on clinic attendances, hospital admissions, and child mortality. Ghana VAST Study Team. Lancet, 342, 7-12.
- Anses/CIQUAL (Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail/Centre d'information sur la qualité des aliments), 2012. French food composition table version 2012. Available online: <http://www.afssa.fr/TableCIQUAL/index.htm>.
- Arnhold T, Tzimas G, Wittfoht W, Plonait S and Nau H, 1996. Identification of 9-cis-retinoic acid, 9,13-di-cis-retinoic acid, and 14-hydroxy-4,14-retro-retinol in human plasma after liver consumption. Life Sciences, 59, 169-177.
- Asgari MM, Brasky TM and White E, 2012. Association of vitamin A and carotenoid intake with melanoma risk in a large prospective cohort. Journal of Investigative Dermatology, 132, 1573-1582.
- Ballev C, Bowman BA, Sowell AL and Gillespie C, 2001. Serum retinol distributions in residents of the United States: third National Health and Nutrition Examination Survey, 1988-1994. American Journal of Clinical Nutrition, 73, 586-593.
- Baly DL, Golub MS, Gershwin ME and Hurley LS, 1984. Studies of marginal zinc deprivation in rhesus monkeys. III. Effects on vitamin A metabolism. American Journal of Clinical Nutrition, 40, 199-207.
- Bankson DD, Russell RM and Sadowski JA, 1986. Determination of retinyl esters and retinol in serum or plasma by normal-phase liquid chromatography: method and applications. Clinical Chemistry, 32, 35-40.
- Bankson DD, Ellis JK and Russell RM, 1989. Effects of a vitamin-A-free diet on tissue vitamin A concentration and dark adaptation of aging rats. Experimental Gerontology, 24, 127-136.
- Barua AB and Olson JA, 1986. Retinoyl beta-glucuronide: an endogenous compound of human blood. American Journal of Clinical Nutrition, 43, 481-485.
- Barua AB, Batres RO and Olson JA, 1989. Characterization of retinyl beta-glucuronide in human blood. American Journal of Clinical Nutrition, 50, 370-374.
- Bates B, Lennox A, Prentice A, Bates CJ and Swan G, 2012. Headline results from Years 1,2 and 3 (combined) of the Rolling Programme (2008/2009-2010/11) Department of Health and Food Standards Agency, 79 pp.
- Bauernfeind JC, 1972. Carotenoid vitamin A precursors and analogs in foods and feeds. Journal of Agricultural and Food Chemistry, 20, 456-473.
- Bausch J and Rietz P, 1977. Method for the assessment of vitamin A liver stores. Acta Vitaminologica et Enzymologica, 31, 99-112.
- Beaton GH, Martorell KJ, Aronson KJ, Edmonston B, McCabe G, Ross AC and Harvey B, 1993. Effectiveness of vitamin A supplementation in the control of young child morbidity and mortality in developing countries. Nutrition Policy Discussion Paper No. 13. ACC/SCN State-of-the-art Series. 162 pp.
- Berry DC, Jacobs H, Marwarha G, Gely-Pernot A, O'Byrne SM, DeSantis D, Klopfenstein M, Feret B, Denefeld C, Blaner WS, Croniger CM, Mark M, Noy N and Ghyselinck NB, 2013. The STRA6 receptor is essential for retinol-binding protein-induced insulin resistance but not for maintaining vitamin A homeostasis in tissues other than the eye. Journal of Biological Chemistry, 288, 24528-24539.

- Beydoun MA, Canas JA, Beydoun HA, Chen X, Shroff MR and Zonderman AB, 2012. Serum antioxidant concentrations and metabolic syndrome are associated among U.S. adolescents in recent national surveys. *Journal of Nutrition*, 142, 1693-1704.
- Biesalski HK, 1989. Comparative assessment of the toxicology of vitamin A and retinoids in man. *Toxicology*, 57, 117-161.
- Bjelakovic G, Nagorni A, Nikolova D, Simonetti RG, Bjelakovic M and Gluud C, 2006. Meta-analysis: antioxidant supplements for primary and secondary prevention of colorectal adenoma. *Alimentary Pharmacology and Therapeutics*, 24, 281-291.
- Bjelakovic G, Nikolova D, Simonetti RG and Gluud C, 2008. Systematic review: primary and secondary prevention of gastrointestinal cancers with antioxidant supplements. *Alimentary Pharmacology and Therapeutics*, 28, 689-703.
- Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG and Gluud C, 2012. Antioxidant supplements for prevention of mortality in healthy participants and patients with various diseases. *Cochrane Database of Systematic Reviews*, 3, CD007176.
- Bjelakovic G, Nikolova D and Gluud C, 2013. Meta-regression analyses, meta-analyses, and trial sequential analyses of the effects of supplementation with Beta-carotene, vitamin a, and vitamin e singly or in different combinations on all-cause mortality: do we have evidence for lack of harm? *PLoS ONE*, 8, e74558.
- Bjelakovic G, Nikolova D and Gluud C, 2014. Antioxidant supplements and mortality. *Current Opinion in Clinical Nutrition and Metabolic Care*, 17, 40-44.
- Blaner WS, Hendriks HF, Brouwer A, de Leeuw AM, Knook DL and Goodman DS, 1985. Retinoids, retinoid-binding proteins, and retinyl palmitate hydrolase distributions in different types of rat liver cells. *Journal of Lipid Research*, 26, 1241-1251.
- Blaner WS, Obunike JC, Kurlandsky SB, al-Haideri M, Piantedosi R, Deckelbaum RJ and Goldberg IJ, 1994. Lipoprotein lipase hydrolysis of retinyl ester. Possible implications for retinoid uptake by cells. *Journal of Biological Chemistry*, 269, 16559-16565.
- Blaner WS, 2007. STRA6, a cell-surface receptor for retinol-binding protein: the plot thickens. *Cell Metabolism*, 5, 164-166.
- Blaner WS, O'Byrne SM, Wongsiriroj N, Kluwe J, D'Ambrosio DM, Jiang H, Schwabe RF, Hillman EM, Piantedosi R and Libien J, 2009. Hepatic stellate cell lipid droplets: a specialized lipid droplet for retinoid storage. *Biochimica et Biophysica Acta: Protein Structure and Molecular Enzymology*, 1791, 467-473.
- Bloem MW, Wedel M, Egger RJ, Speek AJ, Schrijver J, Saowakontha S and Schreurs WH, 1989. Iron metabolism and vitamin A deficiency in children in northeast Thailand. *American Journal of Clinical Nutrition*, 50, 332-338.
- Blomhoff R, Green MH, Berg T and Norum KR, 1990. Transport and storage of vitamin A. *Science*, 250, 399-404.
- Blomhoff R, Green MH, Green JB, Berg T and Norum KR, 1991. Vitamin A metabolism: new perspectives on absorption, transport, and storage. *Physiological Reviews*, 71, 951-990.
- Blomhoff R and Blomhoff HK, 2006. Overview of retinoid metabolism and function. *Journal of Neurobiology*, 66, 606-630.
- Borel P, Dubois C, Mekki N, Grolier P, Partier A, Alexandre-Gouabau MC, Lairon D and Azais-Braesco V, 1997. Dietary triglycerides, up to 40 g/meal, do not affect preformed vitamin A bioavailability in humans. *European Journal of Clinical Nutrition*, 51, 717-722.
- Borel P, Moussa M, Reboul E, Lyan B, Defoort C, Vincent-Baudry S, Maillot M, Gastaldi M, Darmon M, Portugal H, Planells R and Lairon D, 2007. Human plasma levels of vitamin E and carotenoids

- are associated with genetic polymorphisms in genes involved in lipid metabolism. *Journal of Nutrition*, 137, 2653-2659.
- Bouillet P, Sapin V, Chazaud C, Messaddeq N, Decimo D, Dolle P and Chambon P, 1997. Developmental expression pattern of *Stra6*, a retinoic acid-responsive gene encoding a new type of membrane protein. *Mechanisms of Development*, 63, 173-186.
- Brown N and Roberts C, 2004. Vitamin A for acute respiratory infection in developing countries: a meta-analysis. *Acta Paediatrica*, 93, 1437-1442.
- Butte NF, Lopez-Alarcon MG and Garza C, 2002. Nutrient adequacy of exclusive breastfeeding for the term infant during the first six months of life. *World Health Organization*, 57 pp.
- Caire-Juvera G, Ritenbaugh C, Wactawski-Wende J, Snetselaar LG and Chen Z, 2009. Vitamin A and retinol intakes and the risk of fractures among participants of the Women's Health Initiative Observational Study. *American Journal of Clinical Nutrition*, 89, 323-330.
- Canfield LM, Clandinin MT, Davies DP, Fernandez MC, Jackson J, Hawkes J, Goldman WJ, Pramuk K, Reyes H, Sablan B, Sonobe T and Bo X, 2003. Multinational study of major breast milk carotenoids of healthy mothers. *European Journal of Nutrition*, 42, 133-141.
- Carney EA and Russell RM, 1980. Correlation of dark adaptation test results with serum vitamin A levels in diseased adults. *Journal of Nutrition*, 110, 552-557.
- Cassani B, Villablanca EJ, De Calisto J, Wang S and Mora JR, 2012. Vitamin A and immune regulation: role of retinoic acid in gut-associated dendritic cell education, immune protection and tolerance. *Molecular Aspects of Medicine*, 33, 63-76.
- Castenmiller JJ and West CE, 1998. Bioavailability and bioconversion of carotenoids. *Annual Review of Nutrition*, 18, 19-38.
- Chasan-Taber L, Willett WC, Seddon JM, Stampfer MJ, Rosner B, Colditz GA, Speizer FE and Hankinson SE, 1999. A prospective study of carotenoid and vitamin A intakes and risk of cataract extraction in US women. *American Journal of Clinical Nutrition*, 70, 509-516.
- Chase HP, Kumar V, Dodds JM, Sauberlich HE, Hunter RM, Burton RS and Spalding V, 1971. Nutritional status of preschool Mexican-American migrant farm children. *American Journal of Diseases of Children*, 122, 316-324.
- Chen H, Zhuo Q, Yuan W, Wang J and Wu T, 2008. Vitamin A for preventing acute lower respiratory tract infections in children up to seven years of age. *Cochrane Database of Systematic Reviews*, 1, 006090.
- Chichili GR, Nohr D, Schaffer M, von Lintig J and Biesalski HK, 2005. beta-Carotene conversion into vitamin A in human retinal pigment epithelial cells. *Investigative Ophthalmology and Visual Science*, 46, 3562-3569.
- Christian P, West KP, Jr., Khattry SK, Katz J, LeClerq S, Pradhan EK and Shrestha SR, 1998a. Vitamin A or beta-carotene supplementation reduces but does not eliminate maternal night blindness in Nepal. *Journal of Nutrition*, 128, 1458-1463.
- Christian P, Schulze K, Stoltzfus RJ and West KP, Jr., 1998b. Hyporetinolemia, illness symptoms, and acute phase protein response in pregnant women with and without night blindness. *American Journal of Clinical Nutrition*, 67, 1237-1243.
- Christian P and West KP, Jr., 1998. Interactions between zinc and vitamin A: an update. *American Journal of Clinical Nutrition*, 68, 435S-441S.
- Chuwers P, Barnhart S, Blanc P, Brodtkin CA, Cullen M, Kelly T, Keogh J, Omenn G, Williams J and Balmes JR, 1997. The protective effect of beta-carotene and retinol on ventilatory function in an asbestos-exposed cohort. *American Journal of Respiratory and Critical Care Medicine*, 155, 1066-1071.

- Cifelli CJ, Green JB, Wang Z, Yin S, Russell RM, Tang G and Green MH, 2008. Kinetic analysis shows that vitamin A disposal rate in humans is positively correlated with vitamin A stores. *Journal of Nutrition*, 138, 971-977.
- Clagett-Dame M and Knutson D, 2011. Vitamin A in reproduction and development. *Nutrients*, 3, 385-428.
- Cohen-Tanugi A and Forest N, 1998. Retinoic acid suppresses the osteogenic differentiation capacity of murine osteoblast-like 3A/1D-1M cell cultures. *Differentiation*, 63, 115-123.
- Cumming RG, Mitchell P and Smith W, 2000. Diet and cataract: the Blue Mountains Eye Study. *Ophthalmology*, 107, 450-456.
- D'Ambrosio DN, Clugston RD and Blaner WS, 2011. Vitamin A metabolism: an update. *Nutrients*, 3, 63-103.
- D-A-CH (Deutsche Gesellschaft für Ernährung - Österreichische Gesellschaft für Ernährung - Schweizerische Gesellschaft für Ernährungsforschung - Schweizerische Vereinigung für Ernährung), 2013. Referenzwerte für die Nährstoffzufuhr. Neuer Umschau Buchverlag, Neustadt an der Weinstraße, Germany, 292 pp.
- Davis HR, Jr., Zhu LJ, Hoos LM, Tetzloff G, Maguire M, Liu J, Yao X, Iyer SP, Lam MH, Lund EG, Detmers PA, Graziano MP and Altmann SW, 2004. Niemann-Pick C1 Like 1 (NPC1L1) is the intestinal phytosterol and cholesterol transporter and a key modulator of whole-body cholesterol homeostasis. *Journal of Biological Chemistry*, 279, 33586-33592.
- de Pee S, West CE, Permaesih D, Martuti S, Muhilal and Hautvast JG, 1998. Orange fruit is more effective than are dark-green, leafy vegetables in increasing serum concentrations of retinol and beta-carotene in schoolchildren in Indonesia. *American Journal of Clinical Nutrition*, 68, 1058-1067.
- de Pee S and Bloem MW, 2007. The bioavailability of (pro) vitamin A carotenoids and maximizing the contribution of homestead food production to combating vitamin A deficiency. *International Journal for Vitamin and Nutrition Research*, 77, 182-192.
- Devery J and Milborrow BV, 1994. beta-Carotene-15,15'-dioxygenase (EC 1.13.11.21) isolation reaction mechanism and an improved assay procedure. *British Journal of Nutrition*, 72, 397-414.
- DH (Department of Health), 1991. Dietary reference values for food energy and nutrients for the United Kingdom. Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy. 212 pp.
- Dorea JG and Olson JA, 1986. The rate of rhodopsin regeneration in the bleached eyes of zinc-deficient rats in the dark. *Journal of Nutrition*, 116, 121-127.
- Dowling JE and Gibbons IR, 1961. The effect of vitamin A deficiency of the fine structure of the retina. In: *The Structure of the Eye*. Ed Smelser GK. Academic Press, New York, USA pp. 85-99.
- Duda G, Nogala-Kalucka M, Karwowska W, Kupczyk B and Lampart-Szczapa E, 2009. Influence of the lactating women diet on the concentration of the lipophilic vitamins in human milk. *Pakistan Journal of Nutrition*, 8, 629-634.
- Duncan JR and Hurley LS, 1978. An interaction between zinc and vitamin A in pregnant and fetal rats. *Journal of Nutrition*, 108, 1431-1438.
- During A, Smith MK, Piper JB and Smith JC, 2001. beta-Carotene 15,15'-Dioxygenase activity in human tissues and cells: evidence of an iron dependency. *Journal of Nutritional Biochemistry*, 12, 640-647.
- During A, Hussain MM, Morel DW and Harrison EH, 2002. Carotenoid uptake and secretion by CaCo-2 cells: beta-carotene isomer selectivity and carotenoid interactions. *Journal of Lipid Research*, 43, 1086-1095.

- During A and Harrison EH, 2004. Intestinal absorption and metabolism of carotenoids: insights from cell culture. *Archives of Biochemistry and Biophysics*, 430, 77-88.
- During A, Dawson HD and Harrison EH, 2005. Carotenoid transport is decreased and expression of the lipid transporters SR-BI, NPC1L1, and ABCA1 is downregulated in Caco-2 cells treated with ezetimibe. *Journal of Nutrition*, 135, 2305-2312.
- During A and Harrison EH, 2007. Mechanisms of provitamin A (carotenoid) and vitamin A (retinol) transport into and out of intestinal Caco-2 cells. *Journal of Lipid Research*, 48, 2283-2294.
- EFSA (European Food Safety Authority), 2011a. Report on the development of a food classification and description system for exposure assessment and guidance on its implementation and use. *EFSA Journal* 2011;9(12):2489, 84 pp. doi:10.2903/j.efsa.2011.2489
- EFSA (European Food Safety Authority), 2011b. Use of the EFSA Comprehensive European Food Consumption Database in exposure assessment. *EFSA Journal* 2011;9(3):2097, 34 pp. doi:10.2903/j.efsa.2011.2097
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2009. Scientific Opinion on the appropriate age for introduction of complementary feeding of infants. *EFSA Journal* 2009;7(12):1423, 38 pp. doi:10.2903/j.efsa.2009.1423.
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2013. Scientific Opinion on Dietary Reference Values for energy. *EFSA Journal* 2013;11(1):3005, 112 pp. doi:10.2903/j.efsa.2013.3005
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2014a. Scientific Opinion on Dietary Reference Values for selenium. *EFSA Journal* 2014;12(10):3846, 66 pp. doi:10.2903/j.efsa.2014.3846
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2014b. Scientific Opinion on the essential composition of infant and follow-on formulae. *EFSA Journal* 2014;12(7):3760, 106 pp. doi:10.2903/j.efsa.2014.3760
- Erlanson C and Borgstrom B, 1968. The identity of vitamin A esterase activity of rat pancreatic juice. *Biochimica et Biophysica Acta: Protein Structure and Molecular Enzymology*, 167, 629-631.
- Eroglu A and Harrison EH, 2013. Carotenoid metabolism in mammals, including man: formation, occurrence, and function of apocarotenoids. *Journal of Lipid Research*, 54, 1719-1730.
- Fairfield KM, Hankinson SE, Rosner BA, Hunter DJ, Colditz GA and Willett WC, 2001. Risk of ovarian carcinoma and consumption of vitamins A, C, and E and specific carotenoids: a prospective analysis. *Cancer*, 92, 2318-2326.
- FAO/WHO (Food and Agriculture Organization of the United Nations/World Health Organization), 1988. Requirements of Vitamin A, Iron, Folate and Vitamin B12. Report of a joint FAO/WHO Expert Consultation. Food and Agriculture Organization of the United Nations, Rome, Italy, 107 pp.
- FAO/WHO/UNU (Food and Agriculture Organization of the United Nations/World Health Organization/United Nations University), 2004. Human energy requirements. Report of a Joint FAO/WHO/UNU Expert Consultation: Rome 17-24 October 2001. FAO food and nutrition technical report series, 103 pp.
- Farjo KM, Moiseyev G, Nikolaeva O, Sandell LL, Trainor PA and Ma JX, 2011. RDH10 is the primary enzyme responsible for the first step of embryonic Vitamin A metabolism and retinoic acid synthesis. *Developmental Biology*, 357, 347-355.
- Fawzi WW, Herrera MG and Nestel P, 1992. Vitamin A supplementation in malnourished Sudanese children. *International Journal for Vitamin and Nutrition Research*, 62, 344.
- Ferrucci L, Perry JR, Matteini A, Perola M, Tanaka T, Silander K, Rice N, Melzer D, Murray A, Cluett C, Fried LP, Albanes D, Corsi AM, Cherubini A, Guralnik J, Bandinelli S, Singleton A,

- Virtamo J, Walston J, Semba RD and Frayling TM, 2009. Common variation in the beta-carotene 15,15'-monooxygenase 1 gene affects circulating levels of carotenoids: a genome-wide association study. *American Journal of Human Genetics*, 84, 123-133.
- Feskanich D, Singh V, Willett WC and Colditz GA, 2002. Vitamin A intake and hip fractures among postmenopausal women. *JAMA*, 287, 47-54.
- Feskanich D, Willett WC, Hunter DJ and Colditz GA, 2003. Dietary intakes of vitamins A, C, and E and risk of melanoma in two cohorts of women. *British Journal of Cancer*, 88, 1381-1387.
- Field CJ, Johnson IR and Schley PD, 2002. Nutrients and their role in host resistance to infection. *Journal of Leukocyte Biology*, 71, 16-32.
- Field S, Elliott F, Randerson-Moor J, Kukalich K, Barrett JH, Bishop DT and Newton-Bishop JA, 2013. Do vitamin A serum levels moderate outcome or the protective effect of vitamin D on outcome from malignant melanoma? *Clinical Nutrition*, 32, 1012-1016.
- Figueira F, Mendonca S, Rocha J, Azevedo M, Bunce GE and Reynolds JW, 1969. Absorption of vitamin A by infants receiving fat-free or fat-containing dried skim milk formulas. *American Journal of Clinical Nutrition*, 22, 588-593.
- Filteau SM, Morris SS, Abbott RA, Tomkins AM, Kirkwood BR, Arthur P, Ross DA, Gyapong JO and Raynes JG, 1993. Influence of morbidity on serum retinol of children in a community-based study in northern Ghana. *American Journal of Clinical Nutrition*, 58, 192-197.
- Fletcher AE, Breeze E and Shetty PS, 2003. Antioxidant vitamins and mortality in older persons: findings from the nutrition add-on study to the Medical Research Council Trial of Assessment and Management of Older People in the Community. *American Journal of Clinical Nutrition*, 78, 999-1010.
- Flores H and de Araujo CR, 1984. Liver levels of retinol in unselected necropsy specimens: a prevalence survey of vitamin A deficiency in Recife, Brazil. *American Journal of Clinical Nutrition*, 40, 146-152.
- Flores H, Azevedo MN, Campos FA, Barreto-Lins MC, Cavalcanti AA, Salzano AC, Varela RM and Underwood BA, 1991. Serum vitamin A distribution curve for children aged 2-6 y known to have adequate vitamin A status: a reference population. *American Journal of Clinical Nutrition*, 54, 707-711.
- Flores H, 1993. Frequency distributions of serum vitamin A levels in cross-sectional surveys and in surveys before and after vitamin A supplementation. In: *A brief guide to current methods of assessing vitamin A status. A report of the International Vitamin A Consultative Group (IVACG)*. The Nutrition Foundation, Washington, DC, USA, 9-11.
- Fortmann SP, Burda BU, Senger CA, Lin JS and Whitlock EP, 2013. Vitamin and mineral supplements in the primary prevention of cardiovascular disease and cancer: An updated systematic evidence review for the U.S. Preventive Services Task Force. *Annals of Internal Medicine*, 159, 824-834.
- Freudenheim JL, Johnson NE and Smith EL, 1986. Relationships between usual nutrient intake and bone-mineral content of women 35-65 years of age: longitudinal and cross-sectional analysis. *American Journal of Clinical Nutrition*, 44, 863-876.
- FSA (Food Standards Agency), 2002. McCance and Widdowson's The Composition of Foods integrated dataset. Available online: <http://www.food.gov.uk/science/dietarysurveys/dietsurveys/>.
- Fujii H, Sato T, Kaneko S, Gotoh O, Fujii-Kuriyama Y, Osawa K, Kato S and Hamada H, 1997. Metabolic inactivation of retinoic acid by a novel P450 differentially expressed in developing mouse embryos. *EMBO Journal*, 16, 4163-4173.
- Fujita M, Brindle E, Rocha A, Shell-Duncan B, Ndemwa P and O'Connor KA, 2009. Assessment of the relative dose-response test based on serum retinol-binding protein instead of serum retinol in determining low hepatic vitamin A stores. *American Journal of Clinical Nutrition*, 90, 217-224.

- Fulan H, Changxing J, Baina WY, Wencui Z, Chunqing L, Fan W, Dandan L, Dianjun S, Tong W, Da P and Yashuang Z, 2011. Retinol, vitamins A, C, and E and breast cancer risk: a meta-analysis and meta-regression. *Cancer Causes & Control*, 22, 1383-1396.
- Fung TT, Hunter DJ, Spiegelman D, Colditz GA, Speizer FE and Willett WC, 2002. Vitamins and carotenoids intake and the risk of basal cell carcinoma of the skin in women (United States). *Cancer Causes & Control*, 13, 221-230.
- Furr HC, Amedee-Manesme O, Clifford AJ, Bergen HR, 3rd, Jones AD, Anderson DP and Olson JA, 1989. Vitamin A concentrations in liver determined by isotope dilution assay with tetradeuterated vitamin A and by biopsy in generally healthy adult humans. *American Journal of Clinical Nutrition*, 49, 713-716.
- Furr HC, Green MH, Haskell M, Mokhtar N, Nestel P, Newton S, Ribaya-Mercado JD, Tang G, Tanumihardjo S and Wasantwisut E, 2005. Stable isotope dilution techniques for assessing vitamin A status and bioefficacy of provitamin A carotenoids in humans. *Public health nutrition*, 8, 596-607.
- Gadomski AM, Kjolhede CL, Wittppenn J, Bulux J, Rosas AR and Forman MR, 1989. Conjunctival impression cytology (CIC) to detect subclinical vitamin A deficiency: comparison of CIC with biochemical assessments. *American Journal of Clinical Nutrition*, 49, 495-500.
- Garcia-Closas R, Castellsague X, Bosch X and Gonzalez CA, 2005. The role of diet and nutrition in cervical carcinogenesis: a review of recent evidence. *International Journal of Cancer*, 117, 629-637.
- Garcia-Closas R, Garcia-Closas M, Kogevinas M, Malats N, Silverman D, Serra C, Tardon A, Carrato A, Castano-Vinyals G, Dosemeci M, Moore L, Rothman N and Sinha R, 2007. Food, nutrient and heterocyclic amine intake and the risk of bladder cancer. *European Journal of Cancer*, 43, 1731-1740.
- Ghadirian P, Lacroix A, Maisonneuve P, Perret C, Drouin G, Perrault JP, Beland G, Rohan TE and Howe GR, 1996. Nutritional factors and prostate cancer: a case-control study of French Canadians in Montreal, Canada. *Cancer Causes & Control*, 7, 428-436.
- Giaconi JA, Yu F, Stone KL, Pedula KL, Ensrud KE, Cauley JA, Hochberg MC and Coleman AL, 2012. The association of consumption of fruits/vegetables with decreased risk of glaucoma among older African-American women in the study of osteoporotic fractures. *American Journal of Ophthalmology*, 154, 635-644.
- Giovannucci E, Ascherio A, Rimm EB, Stampfer MJ, Colditz GA and Willett WC, 1995. Intake of carotenoids and retinol in relation to risk of prostate cancer. *Journal of the National Cancer Institute*, 87, 1767-1776.
- Glasziou PP and Mackerras DE, 1993. Vitamin A supplementation in infectious diseases: a meta-analysis. *British Medical Journal*, 306, 366-370.
- Glatz JF, van Nieuwenhoven FA, Luiken JJ, Schaap FG and van der Vusse GJ, 1997. Role of membrane-associated and cytoplasmic fatty acid-binding proteins in cellular fatty acid metabolism. *Prostaglandins Leukotrienes and Essential Fatty Acids*, 57, 373-378.
- Gonzalez CA, Travier N, Lujan-Barroso L, Castellsague X, Bosch FX, Roura E, Bueno-de-Mesquita HB, Palli D, Boeing H, Pala V, Sacerdote C, Tumino R, Panico S, Manjer J, Dillner J, Hallmans G, Kjellberg L, Sanchez MJ, Altzibar JM, Barricarte A, Navarro C, Rodriguez L, Allen N, Key TJ, Kaaks R, Rohrmann S, Overvad K, Olsen A, Tjonneland A, Munk C, Kjaer SK, Peeters PH, van Duijnhoven FJ, Clavel-Chapelon F, Boutron-Ruault MC, Trichopoulou A, Benetou V, Naska A, Lund E, Engeset D, Skeie G, Franceschi S, Slimani N, Rinaldi S and Riboli E, 2011. Dietary factors and in situ and invasive cervical cancer risk in the European prospective investigation into cancer and nutrition study. *International Journal of Cancer*, 129, 449-459.
- Goodman DW, Huang HS and Shiratori T, 1965. Tissue distribution and metabolism of newly absorbed vitamin a in the rat. *Journal of Lipid Research*, 6, 390-396.

- Green MH and Green JB, 1994. Vitamin A intake and status influence retinol balance, utilization and dynamics in rats. *Journal of Nutrition*, 124, 2477-2485.
- Grotto I, Mimouni M, Gdalevich M and Mimouni D, 2003. Vitamin A supplementation and childhood morbidity from diarrhea and respiratory infections: a meta-analysis. *Journal of Pediatrics*, 142, 297-304.
- Haddad S, Restieri C and Krishnan K, 2001. Characterization of age-related changes in body weight and organ weights from birth to adolescence in humans. *Journal of Toxicology and Environmental Health. Part A*, 64, 453-464.
- Harrison EH, Blaner WS, Goodman DS and Ross AC, 1987. Subcellular localization of retinoids, retinoid-binding proteins, and acyl-CoA:retinol acyltransferase in rat liver. *Journal of Lipid Research*, 28, 973-981.
- Harrison EH, 2012. Mechanisms involved in the intestinal absorption of dietary vitamin A and provitamin A carotenoids. *Biochimica et Biophysica Acta: Protein Structure and Molecular Enzymology*, 1821, 70-77.
- Hartmann S, Froescheis O, Ringenbach F, Wyss R, Bucheli F, Bischof S, Bausch J and Wiegand UW, 2001. Determination of retinol and retinyl esters in human plasma by high-performance liquid chromatography with automated column switching and ultraviolet detection. *Journal of Chromatography. B, Biomedical Sciences and Applications*, 751, 265-275.
- Haskell MJ, Handelman GJ, Peerson JM, Jones AD, Rabbi MA, Awal MA, Wahed MA, Mahalanabis D and Brown KH, 1997. Assessment of vitamin A status by the deuterated-retinol-dilution technique and comparison with hepatic vitamin A concentration in Bangladeshi surgical patients. *American Journal of Clinical Nutrition*, 66, 67-74.
- Haskell MJ, Islam MA, Handelman GJ, Peerson JM, Jones AD, Wahed MA, Mahalanabis D and Brown KH, 1998. Plasma kinetics of an oral dose of [2H4]retinyl acetate in human subjects with estimated low or high total body stores of vitamin A. *American Journal of Clinical Nutrition*, 68, 90-95.
- Haskell MJ, Mazumder RN, Peerson JM, Jones AD, Wahed MA, Mahalanabis D and Brown KH, 1999. Use of the deuterated-retinol-dilution technique to assess total-body vitamin A stores of adult volunteers consuming different amounts of vitamin A. *American Journal of Clinical Nutrition*, 70, 874-880.
- Haskell MJ, Lembcke JL, Salazar M, Green MH, Peerson JM and Brown KH, 2003. Population-based plasma kinetics of an oral dose of [2H4]retinyl acetate among preschool-aged, Peruvian children. *American Journal of Clinical Nutrition*, 77, 681-686.
- Haskell MJ, Ribaya-Mercado JD and the Vitamin A Tracer Task Force, 2005. Handbook on Vitamin A tracer dilution methods to assess status and evaluate intervention programs. HarvestPlus Technical Monograph 5, 24 pp.
- Haskell MJ, Jamil KM, Peerson JM, Wahed MA and Brown KH, 2011. The paired deuterated retinol dilution technique can be used to estimate the daily vitamin A intake required to maintain a targeted whole body vitamin A pool size in men. *Journal of Nutrition*, 141, 428-432.
- Haskell MJ, 2012. The challenge to reach nutritional adequacy for vitamin A: beta-carotene bioavailability and conversion--evidence in humans. *American Journal of Clinical Nutrition*, 96, 1193S-1203S.
- Hathcock JN, Hattan DG, Jenkins MY, McDonald JT, Sundaresan PR and Wilkening VL, 1990. Evaluation of vitamin A toxicity. *American Journal of Clinical Nutrition*, 52, 183-202.
- Heinonen M, Kärkkäinen M, Riuttamäki M-A, Piironen V, Lampi A-M, Ollilainen V and Lamberg-Allardt C, 2012. Literature search and review related to specific preparatory work in the establishment of Dietary Reference Values. Preparation of an evidence report identifying health

- outcomes upon which Dietary Reference Values could potentially be based for vitamins A, C, E, and K. EFSA supporting publication 2012:EN-256, 331 pp.
- Helldán A, Raulio S, Kosola M, Tapanainen H, Ovaskainen ML and Virtanen S, 2013. Finravinto 2012 -tutkimus - The National FINDIET 2012 Survey. THL. Raportti 16/2013. Helsinki, Finland, 217 pp.
- Hendriks HF, Verhoofstad WA, Brouwer A, de Leeuw AM and Knook DL, 1985. Perisinusoidal fat-storing cells are the main vitamin A storage sites in rat liver. *Experimental Cell Research*, 160, 138-149.
- Herberg S, Galan P, Preziosi P, Bertrais S, Mennen L, Malvy D, Roussel AM, Favier A and Briancon S, 2004. The SU.VI.MAX Study: a randomized, placebo-controlled trial of the health effects of antioxidant vitamins and minerals. *Archives of Internal Medicine*, 164, 2335-2342.
- Herr FM and Ong DE, 1992. Differential interaction of lecithin-retinol acyltransferase with cellular retinol binding proteins. *Biochemistry*, 31, 6748-6755.
- Heseker H, Schneider R, Moch K, Kohlmeier M and Kübler W, 1994. Vitaminversorgung Erwachsener in der Bundesrepublik Deutschland. VERA-Schriftenreihe Bd. IV. Wissenschaftl. Fachverlag. Fleck, Niederkleen, Germany, 209 pp.
- Hicks VA, Gunning DB and Olson JA, 1984. Metabolism, plasma transport and biliary excretion of radioactive vitamin A and its metabolites as a function of liver reserves of vitamin A in the rat. *Journal of Nutrition*, 114, 1327-1333.
- Ho CC, de Moura FF, Kim SH, Burri BJ and Clifford AJ, 2009. A minute dose of ¹⁴C- β -carotene is absorbed and converted to retinoids in humans. *Journal of Nutrition*, 139, 1480-1486.
- Hollander D and Muralidhara KS, 1977. Vitamin A₁ intestinal absorption in vivo: influence of luminal factors on transport. *American Journal of Physiology*, 232, E471-477.
- Hollander D, 1981. Intestinal absorption of vitamins A, E, D, and K. *Journal of Laboratory and Clinical Medicine*, 97, 449-462.
- Hoppner K, Phillips WE, Erdody P, Murray TK and Perrin DE, 1969. Vitamin A reserves of Canadians. *Canadian Medical Association Journal*, 101, 84-86.
- Houtkooper LB, Ritenbaugh C, Aickin M, Lohman TG, Going SB, Weber JL, Greaves KA, Boyden TW, Pamerter RW and Hall MC, 1995. Nutrients, body composition and exercise are related to change in bone mineral density in premenopausal women. *Journal of Nutrition*, 125, 1229-1237.
- Hsu WL, Pan WH, Chien YC, Yu KJ, Cheng YJ, Chen JY, Liu MY, Hsu MM, Lou PJ, Chen IH, Yang CS, Hildesheim A and Chen CJ, 2012. Lowered risk of nasopharyngeal carcinoma and intake of plant vitamin, fresh fish, green tea and coffee: a case-control study in Taiwan. *PLoS ONE*, 7, 27.
- Huang HS and Goodman DS, 1965. Vitamin a and Carotenoids. I. Intestinal Absorption and Metabolism of ¹⁴c-Labelled Vitamin a Alcohol and Beta-Carotene in the Rat. *Journal of Biological Chemistry*, 240, 2839-2844.
- Hume EM and Krebs HA, 1949. Vitamin A requirements of human adults. In: Report of the vitamin A subcommittee of the Accessory Food Factors Committee. Special Report series N°264. Eds Hume EM and Krebs HA. Medical Research Council, HM Stationery Office, London, UK, 7-145.
- Huque T, 1982. A survey of human liver reserves of retinol in London. *British Journal of Nutrition*, 47, 165-172.
- IAEA (International Atomic Energy Agency), 2008. Analysis of stable isotope data to estimate vitamin A body stores. Jointly prepared by HarvestPlus, IAEA and USAID. 27 pp.
- Imdad A, Yakoob MY, Sudfeld C, Haider BA, Black RE and Bhutta ZA, 2011. Impact of vitamin A supplementation on infant and childhood mortality. *BMC Public Health*, 11 Suppl 3, S20.

- IOM (Institute of Medicine), 2001. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. National Academy Press, Washington, DC, USA. 773 pp.
- Iqbal J and Hussain MM, 2009. Intestinal lipid absorption. *American Journal of Physiology, Endocrinology and Metabolism*, 296, E1183-1194.
- IUNA (Irish Universities Nutrition Alliance), 2011. National Adult Nutrition Survey. 40 pp.
- Jalal F, Nesheim MC, Agus Z, Sanjur D and Habicht JP, 1998. Serum retinol concentrations in children are affected by food sources of beta-carotene, fat intake, and anthelmintic drug treatment. *American Journal of Clinical Nutrition*, 68, 623-629.
- Jansen RJ, Robinson DP, Stolzenberg-Solomon RZ, Bamlet WR, de Andrade M, Oberg AL, Rabe KG, Anderson KE, Olson JE, Sinha R and Petersen GM, 2013. Nutrients from fruit and vegetable consumption reduce the risk of pancreatic cancer. *Journal of Gastrointestinal Cancer*, 44, 152-161.
- Jayarajan P, Reddy V and Mohanram M, 1980. Effect of dietary fat on absorption of beta carotene from green leafy vegetables in children. *Indian Journal of Medical Research*, 71, 53-56.
- Johansson S and Melhus H, 2001. Vitamin A antagonizes calcium response to vitamin D in man. *Journal of Bone and Mineral Research*, 16, 1899-1905.
- Johnson EJ and Russell RM, 1992. Distribution of orally administered beta-carotene among lipoproteins in healthy men. *American Journal of Clinical Nutrition*, 56, 128-135.
- Kabat GC, Kim MY, Wactawski-Wende J, Shikany JM, Vitolins MZ and Rohan TE, 2012. Intake of antioxidant nutrients and risk of non-Hodgkin's Lymphoma in the Women's Health Initiative. *Nutrition and Cancer*, 64, 245-254.
- Kane JP and Havel RJ, 1995. Disorders of the biogenesis and secretion of lipoproteins containing the B apolipoproteins. In: *The Metabolic and Molecular Bases of Inherited Disorders*. Eds Scriver CR, Beaudet AL, Sly WS and Valle D. New York, USA, 1853-1885.
- Kang JH, Pasquale LR, Willett W, Rosner B, Egan KM, Faberowski N and Hankinson SE, 2003. Antioxidant intake and primary open-angle glaucoma: a prospective study. *American Journal of Epidemiology*, 158, 337-346.
- Kasparova M, Plisek J, Solichova D, Krcmova L, Kucerova B, Hronek M and Solich P, 2012. Rapid sample preparation procedure for determination of retinol and alpha-tocopherol in human breast milk. *Talanta*, 93, 147-152.
- Katz J, West KP, Jr., Khattry SK, Thapa MD, LeClerq SC, Pradhan EK, Pokhrel RP and Sommer A, 1995. Impact of vitamin A supplementation on prevalence and incidence of xerophthalmia in Nepal. *Investigative Ophthalmology and Visual Science*, 36, 2577-2583.
- Kawaguchi R, Yu J, Honda J, Hu J, Whitelegge J, Ping P, Wiita P, Bok D and Sun H, 2007. A membrane receptor for retinol binding protein mediates cellular uptake of vitamin A. *Science*, 315, 820-825.
- Kawahara TN, Krueger DC, Engelke JA, Harke JM and Binkley NC, 2002. Short-term vitamin A supplementation does not affect bone turnover in men. *Journal of Nutrition*, 132, 1169-1172.
- Kedishvili NY, 2013. Enzymology of retinoic acid biosynthesis and degradation. *Journal of Lipid Research*, 54, 1744-1760.
- Kersting M and Clausen K, 2003. Ernährungsphysiologische Auswertung einer repräsentativen Verzehrsstudie bei Säuglingen und Kleinkindern VELS mit dem Instrumentarium der DONALD Studie. *Forschungsinstitut für Kinderernährung, Dortmund, Germany*, 103 pp.
- Key TJ, Appleby PN, Masset G, Brunner EJ, Cade JE, Greenwood DC, Stephen AM, Kuh D, Bhaniani A, Powell N and Khaw KT, 2012. Vitamins, minerals, essential fatty acids and colorectal cancer risk in the United Kingdom Dietary Cohort Consortium. *International Journal of Cancer*, 131, 18.

- Khan NC, West CE, de Pee S, Bosch D, Phuong HD, Hulshof PJ, Khoi HH, Verhoef H and Hautvast JG, 2007. The contribution of plant foods to the vitamin A supply of lactating women in Vietnam: a randomized controlled trial. *American Journal of Clinical Nutrition*, 85, 1112-1120.
- Kindmark A, Melhus H, Ljunghall S and Ljunggren O, 1995. Inhibitory effects of 9-cis and all-trans retinoic acid on 1,25(OH)₂ vitamin D₃-induced bone resorption. *Calcified Tissue International*, 57, 242-244.
- Kurlandsky SB, Gamble MV, Ramakrishnan R and Blaner WS, 1995. Plasma delivery of retinoic acid to tissues in the rat. *Journal of Biological Chemistry*, 270, 17850-17857.
- Kushi LH, Folsom AR, Prineas RJ, Mink PJ, Wu Y and Bostick RM, 1996. Dietary antioxidant vitamins and death from coronary heart disease in postmenopausal women. *New England Journal of Medicine*, 334, 1156-1162.
- Kusin JA, Reddy V and Sivakumar B, 1974. Vitamin E supplements and the absorption of a massive dose of vitamin A. *American Journal of Clinical Nutrition*, 27, 774-776.
- Larsson SC, Bergkvist L, Naslund I, Rutegard J and Wolk A, 2007. Vitamin A, retinol, and carotenoids and the risk of gastric cancer: a prospective cohort study. *American Journal of Clinical Nutrition*, 85, 497-503.
- LASER Analytica, 2014. Comprehensive literature search and review of breast milk composition as preparatory work for the setting of dietary reference values for vitamins and minerals. EFSA supporting publication 2014:EN-629, 154 pp.
- Leenders M, Leufkens AM, Siersema PD, van Duijnhoven FJ, Vrieling A, Hulshof PJ, van Gils CH, Overvad K, Roswall N, Kyro C, Boutron-Ruault MC, Fagerhazzi G, Cadeau C, Kuhn T, Johnson T, Boeing H, Aleksandrova K, Trichopoulou A, Klinaki E, Androulidaki A, Palli D, Grioni S, Sacerdote C, Tumino R, Panico S, Bakker MF, Skeie G, Weiderpass E, Jakszyn P, Barricarte A, Maria Huerta J, Molina-Montes E, Arguelles M, Johansson I, Ljuslinder I, Key TJ, Bradbury KE, Khaw KT, Wareham NJ, Ferrari P, Duarte-Salles T, Jenab M, Gunter MJ, Vergnaud AC, Wark PA and Bueno-de-Mesquita HB, 2014. Plasma and dietary carotenoids and vitamins A, C and E and risk of colon and rectal cancer in the European Prospective Investigation into Cancer and Nutrition. *International Journal of Cancer*, 135, 2930-2939.
- Lemieux S, Fontani R, Uffelman KD, Lewis GF and Steiner G, 1998. Apolipoprotein B-48 and retinyl palmitate are not equivalent markers of postprandial intestinal lipoproteins. *Journal of Lipid Research*, 39, 1964-1971.
- Leung WC, Hessel S, Meplan C, Flint J, Oberhauser V, Tourniaire F, Hesketh JE, von Lintig J and Lietz G, 2009. Two common single nucleotide polymorphisms in the gene encoding beta-carotene 15,15'-monooxygenase alter beta-carotene metabolism in female volunteers. *FASEB Journal*, 23, 1041-1053.
- Lewis GF, Carpentier A, Adeli K and Giacca A, 2002. Disordered fat storage and mobilization in the pathogenesis of insulin resistance and type 2 diabetes. *Endocrine Reviews*, 23, 201-229.
- Li E and Norris AW, 1996. Structure/function of cytoplasmic vitamin A-binding proteins. *Annual Review of Nutrition*, 16, 205-234.
- Li E and Tso P, 2003. Vitamin A uptake from foods. *Current Opinion in Lipidology*, 14, 241-247.
- Lim LS, Harnack LJ, Lazovich D and Folsom AR, 2004. Vitamin A intake and the risk of hip fracture in postmenopausal women: the Iowa Women's Health Study. *Osteoporosis International*, 15, 552-559.
- Lindqvist A and Andersson S, 2002. Biochemical properties of purified recombinant human beta-carotene 15,15'-monooxygenase. *Journal of Biological Chemistry*, 277, 23942-23948.
- Lindqvist A, He YG and Andersson S, 2005. Cell type-specific expression of beta-carotene 9',10'-monooxygenase in human tissues. *Journal of Histochemistry and Cytochemistry*, 53, 1403-1412.

- Lindqvist A, Sharvill J, Sharvill DE and Andersson S, 2007. Loss-of-function mutation in carotenoid 15,15'-monooxygenase identified in a patient with hypercarotenemia and hypovitaminosis A. *Journal of Nutrition*, 137, 2346-2350.
- Loerch JD, Underwood BA and Lewis KC, 1979. Response of plasma levels of vitamin A to a dose of vitamin A as an indicator of hepatic vitamin A reserves in rats. *Journal of Nutrition*, 109, 778-786.
- Lynch SR, 1997. Interaction of iron with other nutrients. *Nutrition Reviews*, 55, 102-110.
- Macdonald HM, New SA, Golden MH, Campbell MK and Reid DM, 2004. Nutritional associations with bone loss during the menopausal transition: evidence of a beneficial effect of calcium, alcohol, and fruit and vegetable nutrients and of a detrimental effect of fatty acids. *American Journal of Clinical Nutrition*, 79, 155-165.
- MacDonald PN and Ong DE, 1988. Evidence for a lecithin-retinol acyltransferase activity in the rat small intestine. *Journal of Biological Chemistry*, 263, 12478-12482.
- Macias C and Schweigert FJ, 2001. Changes in the concentration of carotenoids, vitamin A, alpha-tocopherol and total lipids in human milk throughout early lactation. *Annals of Nutrition and Metabolism*, 45, 82-85.
- Marlhens F, Bareil C, Griffoin JM, Zrenner E, Amalric P, Eliaou C, Liu SY, Harris E, Redmond TM, Arnaud B, Claustres M and Hamel CP, 1997. Mutations in RPE65 cause Leber's congenital amaurosis. *Nature Genetics*, 17, 139-141.
- Maserejian NN, Giovannucci E, Rosner B and Joshipura K, 2007. Prospective study of vitamins C, E, and A and carotenoids and risk of oral premalignant lesions in men. *International Journal of Cancer*, 120, 970-977.
- Maslova E, Hansen S, Strom M, Halldorsson TI and Olsen SF, 2014. Maternal intake of vitamins A, E and K in pregnancy and child allergic disease: a longitudinal study from the Danish National Birth Cohort. *British Journal of Nutrition*, 111, 1096-1108.
- Matsuura T and Ross AC, 1993. Regulation of hepatic lecithin: retinol acyltransferase activity by retinoic acid. *Archives of Biochemistry and Biophysics*, 301, 221-227.
- Mayo-Wilson E, Imdad A, Herzer K, Yakoob MY and Bhutta ZA, 2011. Vitamin A supplements for preventing mortality, illness, and blindness in children aged under 5: systematic review and meta-analysis. *British Medical Journal (Clinical Research Edition)*, 343, d5094.
- McLaren DS and Kraemer K, 2012. Mortality and morbidity, especially in relation to infections. *World Review of Nutrition and Dietetics*, 103, 76-92.
- Melhus H, Michaelsson K, Kindmark A, Bergstrom R, Holmberg L, Mallmin H, Wolk A and Ljunghall S, 1998. Excessive dietary intake of vitamin A is associated with reduced bone mineral density and increased risk for hip fracture. *Annals of Internal Medicine*, 129, 770-778.
- Mensink GB, Heseke H, Richter A, Stahl A, Vohmann C, Fischer J, Kohler S and Six J (Robert Koch-Institut & Universität Paderborn), 2007. Ernährungsstudie als KiGGS-Modul (EsKiMo). 143 pp.
- Michaelsson K, Lithell H, Vessby B and Melhus H, 2003. Serum retinol levels and the risk of fracture. *New England Journal of Medicine*, 348, 287-294.
- Mikhak B, Bracci PM and Gong Z, 2012. Intake of vitamins D and A and calcium and risk of non-Hodgkin lymphoma: San Francisco Bay Area population-based case-control study. *Nutrition and Cancer*, 64, 674-684.
- Misotti AM and Gnagnarella P, 2013. Vitamin supplement consumption and breast cancer risk: a review. *Ecancermedicalscience*, 7, 365.
- Mitchell GV, Young M and Seward CR, 1973. Vitamin A and carotene levels of a selected population in metropolitan Washington, D. C. *American Journal of Clinical Nutrition*, 26, 992-997.

- Miyazaki M, Doi Y, Ikeda F, Ninomiya T, Hata J, Uchida K, Shirota T, Matsumoto T, Iida M and Kiyohara Y, 2012. Dietary vitamin A intake and incidence of gastric cancer in a general Japanese population: the Hisayama Study. *Gastric Cancer*, 15, 162-169.
- Money DF, 1978. Vitamin E, selenium, iron, and vitamin A content of livers from Sudden Infant Death Syndrome cases and control children: Interrelationships and possible significance. *New Zealand Journal of Science*, 21, 41-45.
- Montreewasuwat N and Olson JA, 1979. Serum and liver concentrations of vitamin A in Thai fetuses as a function of gestational age. *American Journal of Clinical Nutrition*, 32, 601-606.
- Moriwaki H, Blaner WS, Piantedosi R and Goodman DS, 1988. Effects of dietary retinoid and triglyceride on the lipid composition of rat liver stellate cells and stellate cell lipid droplets. *Journal of Lipid Research*, 29, 1523-1534.
- Morrison SA, Russell RM, Carney EA and Oaks EV, 1978. Zinc deficiency: a cause of abnormal dark adaptation in cirrhotics. *American Journal of Clinical Nutrition*, 31, 276-281.
- Moussa M, Landrier JF, Reboul E, Ghiringhelli O, Comera C, Collet X, Frohlich K, Bohm V and Borel P, 2008. Lycopene absorption in human intestinal cells and in mice involves scavenger receptor class B type I but not Niemann-Pick C1-like 1. *Journal of Nutrition*, 138, 1432-1436.
- Muto Y, Smith JE, Milch PO and Goodman DS, 1972. Regulation of retinol-binding protein metabolism by vitamin A status in the rat. *Journal of Biological Chemistry*, 247, 2542-2550.
- Mwanri L, Worsley A, Ryan P and Masika J, 2000. Supplemental vitamin A improves anemia and growth in anemic school children in Tanzania. *Journal of Nutrition*, 130, 2691-2696.
- Nagao A, Doring A, Hoshino C, Terao J and Olson JA, 1996. Stoichiometric conversion of all trans-beta-carotene to retinal by pig intestinal extract. *Archives of Biochemistry and Biophysics*, 328, 57-63.
- Naldi L, Gallus S, Tavani A, Imberti GL, La Vecchia C and Oncology Study Group of the Italian Group for Epidemiologic Research in D, 2004. Risk of melanoma and vitamin A, coffee and alcohol: a case-control study from Italy. *European Journal of Cancer Prevention*, 13, 503-508.
- Napoli JL, 2012. Physiological insights into all-trans-retinoic acid biosynthesis. *Biochimica et Biophysica Acta*, 1821, 152-167.
- Nayak N, Harrison EH and Hussain MM, 2001. Retinyl ester secretion by intestinal cells: a specific and regulated process dependent on assembly and secretion of chylomicrons. *Journal of Lipid Research*, 42, 272-280.
- Netherlands Food and Nutrition Council, 1992. Recommended dietary allowances 1989 in the Netherlands. 115 pp.
- Newcomer ME, Jamison RS and Ong DE, 1998. Structure and function of retinoid-binding proteins. *Sub-Cellular Biochemistry*, 30, 53-80.
- Nieland TJ, Chroni A, Fitzgerald ML, Maliga Z, Zannis VI, Kirchhausen T and Krieger M, 2004. Cross-inhibition of SR-BI- and ABCA1-mediated cholesterol transport by the small molecules BLT-4 and glyburide. *Journal of Lipid Research*, 45, 1256-1265.
- Nordic Council of Ministers (Nordic Council of Ministers), 2014. Nordic Nutrition Recommendations 2012. Integrating nutrition and physical activity. 627 pp.
- Novotny JA, Dueker SR, Zech LA and Clifford AJ, 1995. Compartmental analysis of the dynamics of beta-carotene metabolism in an adult volunteer. *Journal of Lipid Research*, 36, 1825-1838.
- Noy N and Blaner WS, 1991. Interactions of retinol with binding proteins: studies with rat cellular retinol-binding protein and with rat retinol-binding protein. *Biochemistry*, 30, 6380-6386.
- Noy N, 1992a. The ionization behavior of retinoic acid in aqueous environments and bound to serum albumin. *Biochimica et Biophysica Acta*, 1106, 151-158.

- Noy N, 1992b. The ionization behavior of retinoic acid in lipid bilayers and in membranes. *Biochimica et Biophysica Acta*, 1106, 159-164.
- Nurmatov U, Devereux G and Sheikh A, 2011. Nutrients and foods for the primary prevention of asthma and allergy: systematic review and meta-analysis. *Journal of Allergy and Clinical Immunology*, 127, 724-733 e721-730.
- O'Byrne SM, Wongsiriroj N, Libien J, Vogel S, Goldberg IJ, Baehr W, Palczewski K and Blaner WS, 2005. Retinoid absorption and storage is impaired in mice lacking lecithin:retinol acyltransferase (LRAT). *Journal of Biological Chemistry*, 280, 35647-35657.
- O'Byrne SM and Blaner WS, 2013. Retinol and retinyl esters: biochemistry and physiology. *Journal of Lipid Research*, 54, 1731-1743.
- Olson JA, 1972. The prevention of childhood blindness by the administration of massive doses of vitamin A. *Israel Journal of Medical Sciences*, 8, 1199-1206.
- Olson JA, 1979. Liver vitamin A reserves of neonates, preschool children and adults dying of various causes in Salvador, Brazil. *Archivos Latinoamericanos de Nutricion*, 29, 521-545.
- Olson JA, 1984. Serum levels of vitamin A and carotenoids as reflectors of nutritional status. *Journal of the National Cancer Institute*, 73, 1439-1444.
- Olson JA, 1987. Recommended dietary intakes (RDI) of vitamin A in humans. *American Journal of Clinical Nutrition*, 45, 704-716.
- Olson JA and Hodges RE, 1987. Recommended dietary intakes (RDI) of vitamin C in humans. *American Journal of Clinical Nutrition*, 45, 693-703.
- Olson JA, 1989. Provitamin A function of carotenoids: the conversion of beta-carotene into vitamin A. *Journal of Nutrition*, 119, 105-108.
- Ong DE, 1994. Cellular transport and metabolism of vitamin A: roles of the cellular retinoid-binding proteins. *Nutrition Reviews*, 52, S24-31.
- Opotowsky AR and Bilezikian JP, 2004. Serum vitamin A concentration and the risk of hip fracture among women 50 to 74 years old in the United States: a prospective analysis of the NHANES I follow-up study. *American Journal of Medicine*, 117, 169-174.
- Orhon FS, Ulukol B, Kahya D, Cengiz B, Baskan S and Tezcan S, 2009. The influence of maternal smoking on maternal and newborn oxidant and antioxidant status. *European Journal of Pediatrics*, 168, 975-981.
- Packer L, 2005. Carotenoids and retinoids: molecular aspects and health issues. AOCs Press, Champaign, USA, 350 pp.
- Paik J, Vogel S, Quadro L, Piantedosi R, Gottesman M, Lai K, Hamberger L, Vieira Mde M and Blaner WS, 2004. Vitamin A: overlapping delivery pathways to tissues from the circulation. *Journal of Nutrition*, 134, 276S-280S.
- Palczewski K, 2010. Retinoids for treatment of retinal diseases. *Trends in Pharmacological Sciences*, 31, 284-295.
- Pares X, Farres J, Kedishvili N and Duester G, 2008. Medium- and short-chain dehydrogenase/reductase gene and protein families: Medium-chain and short-chain dehydrogenases/reductases in retinoid metabolism. *Cellular and Molecular Life Sciences*, 65, 3936-3949.
- Parker RS, 1996. Absorption, metabolism, and transport of carotenoids. *FASEB Journal*, 10, 542-551.
- Parker RS, Swanson JE, You CS, Edwards AJ and Huang T, 1999. Bioavailability of carotenoids in human subjects. *Proceedings of the Nutrition Society*, 58, 155-162.
- Pasutto F, Sticht H, Hammersen G, Gillessen-Kaesbach G, Fitzpatrick DR, Nurnberg G, Brasch F, Schirmer-Zimmermann H, Tolmie JL, Chitayat D, Houge G, Fernandez-Martinez L, Keating S,

- Mortier G, Hennekam RC, von der Wense A, Slavotinek A, Meinecke P, Bitoun P, Becker C, Nurnberg P, Reis A and Rauch A, 2007. Mutations in STRA6 cause a broad spectrum of malformations including anophthalmia, congenital heart defects, diaphragmatic hernia, alveolar capillary dysplasia, lung hypoplasia, and mental retardation. *American Journal Of Human Genetics*, 80, 550-560.
- Pennimpede T, Cameron DA, MacLean GA, Li H, Abu-Abed S and Petkovich M, 2010. The role of CYP26 enzymes in defining appropriate retinoic acid exposure during embryogenesis. *Birth Defects Research. Part A, Clinical and Molecular Teratology*, 88, 883-894.
- Pilch SM, 1987. Analysis of vitamin A data from the health and nutrition examination surveys. *Journal of Nutrition*, 117, 636-640.
- Promislow JH, Goodman-Gruen D, Slymen DJ and Barrett-Connor E, 2002. Retinol intake and bone mineral density in the elderly: the Rancho Bernardo Study. *Journal of Bone and Mineral Research*, 17, 1349-1358.
- Quadro L, Blaner WS, Salchow DJ, Vogel S, Piantedosi R, Gouras P, Freeman S, Cosma MP, Colantuoni V and Gottesman ME, 1999. Impaired retinal function and vitamin A availability in mice lacking retinol-binding protein. *EMBO Journal*, 18, 4633-4644.
- Radhika MS, Bhaskaram P, Balakrishna N, Ramalakshmi BA, Devi S and Kumar BS, 2002. Effects of vitamin A deficiency during pregnancy on maternal and child health. *British Journal of Obstetrics and Gynaecology*, 109, 689-693.
- Rahman MM, Mahalanabis D, Wahed MA, Islam M, Habte D, Khaled MA and Alvarez JO, 1995. Conjunctival impression cytology fails to detect subclinical vitamin A deficiency in young children. *Journal of Nutrition*, 125, 1869-1874.
- Rahmathullah L, Underwood BA, Thulasiraj RD, Milton RC, Ramaswamy K, Rahmathullah R and Babu G, 1990. Reduced mortality among children in southern India receiving a small weekly dose of vitamin A. *New England Journal of Medicine*, 323, 929-935.
- Raica N, Jr., Scott J, Lowry L and Sauberlich HE, 1972. Vitamin A concentration in human tissues collected from five areas in the United States. *American Journal of Clinical Nutrition*, 25, 291-296.
- Ramdas WD, Wolfs RC, Kiefte-de Jong JC, Hofman A, de Jong PT, Vingerling JR and Jansonius NM, 2012. Nutrient intake and risk of open-angle glaucoma: the Rotterdam Study. *European Journal of Epidemiology*, 27, 385-393.
- Ray WJ, Bain G, Yao M and Gottlieb DI, 1997. CYP26, a novel mammalian cytochrome P450, is induced by retinoic acid and defines a new family. *Journal of Biological Chemistry*, 272, 18702-18708.
- Reboul E, Berton A, Moussa M, Kreuzer C, Crenon I and Borel P, 2006. Pancreatic lipase and pancreatic lipase-related protein 2, but not pancreatic lipase-related protein 1, hydrolyze retinyl palmitate in physiological conditions. *Biochimica et Biophysica Acta: Protein Structure and Molecular Enzymology*, 1761, 4-10.
- Reddy V and Srikantia SG, 1966. Serum vitamin A in kwashiorkor. *American Journal of Clinical Nutrition*, 18, 105-109.
- Reddy V and Sivakumar B, 1972. Studies on vitamin A absorption in children. *Indian Pediatrics*, 9, 307-310.
- Reddy V (World Health Organisation), 1985. Vitamin A Requirements of Preschool Children. Joint FAO/WHO Expert Group on Requirements for Vitamin A, Iron, Folate and Vitamin B12, Doc. No 6.
- Redlich CA, Grauer JN, Van Bennekum AM, Clever SL, Ponn RB and Blaner WS, 1996. Characterization of carotenoid, vitamin A, and alpha-tocopherol levels in human lung tissue and pulmonary macrophages. *American Journal of Respiratory and Critical Care Medicine*, 154, 1436-1443.

- Redlich CA, Chung JS, Cullen MR, Blaner WS, Van Bennekum AM and Berglund L, 1999. Effect of long-term beta-carotene and vitamin A on serum cholesterol and triglyceride levels among participants in the Carotene and Retinol Efficacy Trial (CARET). *Atherosclerosis*, 145, 425-432.
- Reinersdorff DV, Bush E and Liberato DJ, 1996. Plasma kinetics of vitamin A in humans after a single oral dose of [8,9,19-13C]retinyl palmitate. *Journal of Lipid Research*, 37, 1875-1885.
- Rejnmark L, Vestergaard P, Charles P, Hermann AP, Brot C, Eiken P and Mosekilde L, 2004. No effect of vitamin A intake on bone mineral density and fracture risk in perimenopausal women. *Osteoporosis International*, 15, 872-880.
- Relas H, Gylling H and Miettinen TA, 2000. Effect of stanol ester on postabsorptive squalene and retinyl palmitate. *Metabolism: Clinical and Experimental*, 49, 473-478.
- Ribaya-Mercado JD, Mazariegos M, Tang G, Romero-Abal ME, Mena I, Solomons NW and Russell RM, 1999. Assessment of total body stores of vitamin A in Guatemalan elderly by the deuterated-retinol-dilution method. *American Journal of Clinical Nutrition*, 69, 278-284.
- Ribaya-Mercado JD, Solon FS, Solon MA, Cabal-Barza MA, Perfecto CS, Tang G, Solon JA, Fjeld CR and Russell RM, 2000. Bioconversion of plant carotenoids to vitamin A in Filipino school-aged children varies inversely with vitamin A status. *American Journal of Clinical Nutrition*, 72, 455-465.
- Ribaya-Mercado JD, Solon FS, Fermin LS, Perfecto CS, Solon JA, Dolnikowski GG and Russell RM, 2004. Dietary vitamin A intakes of Filipino elders with adequate or low liver vitamin A concentrations as assessed by the deuterated-retinol-dilution method: implications for dietary requirements. *American Journal of Clinical Nutrition*, 79, 633-641.
- Ribaya-Mercado JD, Maramag CC, Tengco LW, Dolnikowski GG, Blumberg JB and Solon FS, 2007. Carotene-rich plant foods ingested with minimal dietary fat enhance the total-body vitamin A pool size in Filipino schoolchildren as assessed by stable-isotope-dilution methodology. *American Journal of Clinical Nutrition*, 85, 1041-1049.
- Rietz P, Vuilleumier JP, Weber F and Wiss O, 1973. Determination of the vitamin A bodypool of rats by an isotopic dilution method. *Experientia*, 29, 168-170.
- Rietz P, Wiss O and Weber F, 1974. Metabolism of vitamin A and the determination of vitamin A status. *Vitamins and Hormones*, 32, 237-249.
- Rigtrup KM and Ong DE, 1992. A retinyl ester hydrolase activity intrinsic to the brush border membrane of rat small intestine. *Biochemistry*, 31, 2920-2926.
- Rigtrup KM, McEwen LR, Said HM and Ong DE, 1994. Retinyl ester hydrolytic activity associated with human intestinal brush border membranes. *American Journal of Clinical Nutrition*, 60, 111-116.
- Rock CL, Swendseid ME, Jacob RA and McKee RW, 1992. Plasma carotenoid levels in human subjects fed a low carotenoid diet. *Journal of Nutrition*, 122, 96-100.
- Roe MA, Bell S, Oseredczuk M, Christensen T, Westenbrink S, Pakkala H, Presser K and Finglas PM, 2013. Updated food composition database for nutrient intake. Supporting Publications 2013:EN-355, 21 pp.
- Roels OA, Trout M and Dujacquier R, 1958. Carotene balances in boys in Ruanda where vitamin A deficiency is prevalent. *Journal of Nutrition*, 65, 115-127.
- Roels OA, Djaeni S, Trout ME, Lauwtg, Heath A, Poey SH, Tarwotjo MS and Suhadi B, 1963. The effect of protein and fat supplements on vitamin A-deficient Indonesian children. *American Journal of Clinical Nutrition*, 12, 380-387.
- Rohan TE, Cook MG, Potter JD and McMichael AJ, 1990. A case-control study of diet and benign proliferative epithelial disorders of the breast. *Cancer Research*, 50, 3176-3181.

- Rohan TE, Jain M and Miller AB, 1998. A case-cohort study of diet and risk of benign proliferative epithelial disorders of the breast (Canada). *Cancer Causes & Control*, 9, 19-27.
- Rohde CM, Manatt M, Clagett-Dame M and DeLuca HF, 1999. Vitamin A antagonizes the action of vitamin D in rats. *Journal of Nutrition*, 129, 2246-2250.
- Romanchik JE, Morel DW and Harrison EH, 1995. Distributions of carotenoids and alpha-tocopherol among lipoproteins do not change when human plasma is incubated in vitro. *Journal of Nutrition*, 125, 2610-2617.
- Rosales FJ, Jang JT, Pinero DJ, Erikson KM, Beard JL and Ross AC, 1999. Iron deficiency in young rats alters the distribution of vitamin A between plasma and liver and between hepatic retinol and retinyl esters. *Journal of Nutrition*, 129, 1223-1228.
- Ross AC, Chen Q and Ma Y, 2011. Vitamin A and retinoic acid in the regulation of B-cell development and antibody production. *Vitamins and Hormones*, 86, 103-126.
- Ross AC and Zolfaghari R, 2011. Cytochrome P450s in the regulation of cellular retinoic acid metabolism. *Annual Review of Nutrition*, 31, 65-87.
- Ross AC, 2012. Vitamin A and retinoic acid in T cell-related immunity. *American Journal of Clinical Nutrition*, 96, 1166S-1172S.
- Ross AC, 2014. Vitamin A. In: *Modern Nutrition in Health and Disease*. 11th Edition. Eds Ross AC, Caballero B, Cousins RJ, Tucker KL and Ziegler TR. Lippincott Williams & Wilkins, Philadelphia, USA, 260-277.
- Ruder EH, Thiebaut AC, Thompson FE, Potischman N, Subar AF, Park Y, Graubard BI, Hollenbeck AR and Cross AJ, 2011. Adolescent and mid-life diet: risk of colorectal cancer in the NIH-AARP Diet and Health Study. *American Journal of Clinical Nutrition*, 94, 1607-1619.
- Ruhl R, 2007. Effects of dietary retinoids and carotenoids on immune development. *Proceedings of the Nutrition Society*, 66, 458-469.
- SACN (Scientific Advisory Committee on Nutrition), 2005. Review of dietary advice on Vitamin A. pp. 85.
- Sauberlich HE, Hodges RE, Wallace DL, Kolder H, Canham JE, Hood J, Raica N, Jr. and Lowry LK, 1974. Vitamin A metabolism and requirements in the human studied with the use of labeled retinol. *Vitamins and Hormones*, 32, 251-275.
- SCF (Scientific Committee for Food), 1993. Nutrient and energy intakes for the European Community. Reports of the Scientific Committee for Food, 31st Series. Food - Science and Technique, European Commission, Luxembourg. 248 pp.
- SCF (Scientific Committee on Food), 2002. Opinion on the Tolerable Upper Intake Level of preformed vitamin A (retinol and retinyl esters). 16 pp.
- SCF (Scientific Committee on Food), 2003. Report of the Scientific Committee on Food on the revision of essential requirements of infant formulae and follow-on formulae. SCF/CS/NUT/IF/65 Final, 213 pp.
- Scheven BA and Hamilton NJ, 1990. Retinoic acid and 1,25-dihydroxyvitamin D₃ stimulate osteoclast formation by different mechanisms. *Bone*, 11, 53-59.
- Schindler R, Friedrich DH, Kramer M, Wacker HH and Feldheim W, 1988. Size and composition of liver vitamin A reserves of human beings who died of various causes. *International Journal for Vitamin and Nutrition Research*, 58, 146-154.
- Schulz C, Engel U, Kreienberg R and Biesalski HK, 2007. Vitamin A and beta-carotene supply of women with gemini or short birth intervals: a pilot study. *European Journal of Nutrition*, 46, 12-20.

- Schweigert FJ, Bathe K, Chen F, Buscher U and Dudenhausen JW, 2004. Effect of the stage of lactation in humans on carotenoid levels in milk, blood plasma and plasma lipoprotein fractions. *European Journal of Nutrition*, 43, 39-44.
- Seddon JM, Ajani UA, Sperduto RD, Hiller R, Blair N, Burton TC, Farber MD, Gragoudas ES, Haller J and Miller DT, 1994. Dietary carotenoids, vitamins A, C, and E, and advanced age-related macular degeneration. Eye Disease Case-Control Study Group. *JAMA*, 272, 1413-1420.
- Semba RD and Bloem MW, 2002. The anemia of vitamin A deficiency: epidemiology and pathogenesis. *European Journal of Clinical Nutrition*, 56, 271-281.
- Sette S, Le Donne C, Piccinelli R, Arcella D, Turrini A, Leclercq C and Group I-SS, 2011. The third Italian National Food Consumption Survey, INRAN-SCAI 2005-06-part 1: nutrient intakes in Italy. *Nutrition, Metabolism and Cardiovascular Diseases*, 21, 922-932.
- Sivakumar B and Reddy V, 1972. Absorption of labelled vitamin A in children during infection. *British Journal of Nutrition*, 27, 299-304.
- Skare KL and DeLuca HF, 1983. Biliary metabolites of all-trans-retinoic acid in the rat. *Archives of Biochemistry and Biophysics*, 224, 13-18.
- Smith JE, Muto Y, Milch PO and Goodman DS, 1973. The effects of chylomicron vitamin A on the metabolism of retinol-binding protein in the rat. *Journal of Biological Chemistry*, 248, 1544-1549.
- Smith JE, Brown ED and Smith JC, Jr., 1974. The effect of zinc deficiency on the metabolism of retinol-binding protein in the rat. *Journal of Laboratory and Clinical Medicine*, 84, 692-697.
- Smith W, Mitchell P, Webb K and Leeder SR, 1999. Dietary antioxidants and age-related maculopathy: the Blue Mountains Eye Study. *Ophthalmology*, 106, 761-767.
- Solomons NW, Morrow FD, Vasquez A, Bulux J, Guerrero AM and Russell RM, 1990. Test-retest reproducibility of the relative dose response for vitamin A status in Guatemalan adults: issues of diagnostic sensitivity. *Journal of Nutrition*, 120, 738-744.
- Sommer A, 1982. Nutritional factors in corneal xerophthalmia and keratomalacia. *Archives of Ophthalmology*, 100, 399-403.
- Sommer A and West KPJ, 1996. Vitamin A deficiency: health, survival, and vision. Oxford University Press, New York, USA, pp. 464.
- Sommer A, 1982. Nutritional Blindness. Xerophthalmia and Keratomalacia. Oxford University Press, New York, USA, pp. 282.
- Soprano DR and Blaner WS, 1994. Plasma retinol binding protein. In: *The retinoids: biology, chemistry, and medicine*. 2nd ed. Eds Sporn MB, Roberts AB and Goodman DS. Raven Press, New York, USA, 257-281.
- Souci SW, Fachmann W and Kraut H, 2000. *Die Zusammensetzung der Lebensmittel. Nährwert-Tabellen*. Auflage medpharm Scientific Publishers, Stuttgart, Germany, 1008 pp.
- Staab DB, Hodges RE, Metcalf WK and Smith JL, 1984. Relationship between vitamin A and iron in the liver. *Journal of Nutrition*, 114, 840-844.
- Stephensen CB, 2001. Vitamin A, infection, and immune function. *Annual Review of Nutrition*, 21, 167-192.
- Stoltzfus RJ and Underwood BA, 1995. Breast-milk vitamin A as an indicator of the vitamin A status of women and infants. *Bulletin of The World Health Organization*, 73, 703-711.
- Strom K, Gundersen TE, Hansson O, Lucas S, Fernandez C, Blomhoff R and Holm C, 2009. Hormone-sensitive lipase (HSL) is also a retinyl ester hydrolase: evidence from mice lacking HSL. *FASEB Journal*, 23, 2307-2316.

- Suharno D, West CE, Muhilal, Karyadi D and Hautvast JG, 1993. Supplementation with vitamin A and iron for nutritional anaemia in pregnant women in West Java, Indonesia. *Lancet*, 342, 1325-1328.
- Suthutvoravoot S and Olson JA, 1974. Plasma and liver concentration of vitamin A in a normal population of urban Thai. *American Journal of Clinical Nutrition*, 27, 883-891.
- Takata Y, Xiang YB, Yang G, Li H, Gao J, Cai H, Gao YT, Zheng W and Shu XO, 2013. Intakes of fruits, vegetables, and related vitamins and lung cancer risk: results from the Shanghai Men's Health Study (2002-2009). *Nutrition and Cancer*, 65, 51-61.
- Tang G, Qin J, Dolnikowski GG and Russell RM, 2003. Short-term (intestinal) and long-term (postintestinal) conversion of beta-carotene to retinol in adults as assessed by a stable-isotope reference method. *American Journal of Clinical Nutrition*, 78, 259-266.
- Tang G, Hu Y, Yin SA, Wang Y, Dallal GE, Grusak MA and Russell RM, 2012. β -carotene in Golden Rice is as good as β -carotene in oil at providing vitamin A to children. *American Journal of Clinical Nutrition*, 96, 658-664.
- Tanumihardjo SA, 1993. The modified relative dose-response assay. In: *A Brief Guide to Current Methods of Assessing Vitamin A Status. A report of the International Vitamin A Consultative Group (IVACG)*. 14-15.
- Tanumihardjo SA, Muherdiyantiningsih, Permaesih D, Komala, Muhilal, Karyadi D and Olson JA, 1996. Daily supplements of vitamin A (8.4 μ mol, 8000 IU) improve the vitamin A status of lactating Indonesian women. *American Journal of Clinical Nutrition*, 63, 32-35.
- Tanumihardjo SA, 2002. Vitamin A and iron status are improved by vitamin A and iron supplementation in pregnant Indonesian women. *Journal of Nutrition*, 132, 1909-1912.
- Tanumihardjo SA and Penniston KL, 2002. Simplified methodology to determine breast milk retinol concentrations. *Journal of Lipid Research*, 43, 350-355.
- Tanumihardjo SA, Palacios N and Pixley KV, 2010. Provitamin a carotenoid bioavailability: what really matters? *International Journal for Vitamin and Nutrition Research*, 80, 336-350.
- Tanumihardjo SA, 2011. Vitamin A: biomarkers of nutrition for development. *American Journal of Clinical Nutrition*, 94, 658S-665S.
- Terhune MW and Sandstead HH, 1972. Decreased RNA polymerase activity in mammalian zinc deficiency. *Science*, 177, 68-69.
- Thorne-Lyman AL and Fawzi WW, 2012. Vitamin A and carotenoids during pregnancy and maternal, neonatal and infant health outcomes: a systematic review and meta-analysis. *Paediatric and Perinatal Epidemiology*, 26 Suppl 1, 36-54.
- Tokusoglu O, Tansug N, Aksit S, Dinc G, Kasirga E and Ozcan C, 2008. Retinol and (alpha)-tocopherol concentrations in breast milk of Turkish lactating mothers under different socio-economic status. *International Journal of Food Sciences and Nutrition*, 59, 166-174.
- Topletz AR, Thatcher JE, Zelter A, Lutz JD, Tay S, Nelson WL and Isoherranen N, 2012. Comparison of the function and expression of CYP26A1 and CYP26B1, the two retinoic acid hydroxylases. *Biochemical Pharmacology*, 83, 149-163.
- Tsutsumi C, Okuno M, Tannous L, Piantedosi R, Allan M, Goodman DS and Blaner WS, 1992. Retinoids and retinoid-binding protein expression in rat adipocytes. *Journal of Biological Chemistry*, 267, 1805-1810.
- Turley SD and Dietschy JM, 2003. Sterol absorption by the small intestine. *Current Opinion in Lipidology*, 14, 233-240.
- U.S. DH/FAO (US Department of Health, Education, and Welfare/Food and Agriculture Organization), 1968. Food composition table for use in Africa.

- Underwood BA, Siegel H, Weisell RC and Dolinski M, 1970. Liver stores of vitamin A in a normal population dying suddenly or rapidly from unnatural causes in New York City. *American Journal of Clinical Nutrition*, 23, 1037-1042.
- Underwood BA, 1994a. Hypovitaminosis A: international programmatic issues. *Journal of Nutrition*, 124, 1467S-1472S.
- Underwood BA, 1994b. Maternal vitamin A status and its importance in infancy and early childhood. *American Journal of Clinical Nutrition*, 59, 517S-522S.
- Unlu NZ, Bohn T, Clinton SK and Schwartz SJ, 2005. Carotenoid absorption from salad and salsa by humans is enhanced by the addition of avocado or avocado oil. *Journal of Nutrition*, 135, 431-436.
- Valentine AR, 2013. Vitamin A intake requirements in women: Evaluating the Estimated Average Requirement using stable isotope dilution. PhD Thesis, University of Wisconsin-Madison, Madison, United States, 123 pp.
- Valentine AR, Davis CR and Tanumihardjo SA, 2013. Vitamin A isotope dilution predicts liver stores in line with long-term vitamin A intake above the current Recommended Dietary Allowance for young adult women. *American Journal of Clinical Nutrition*, 98, 1192-1199.
- van Bennekum A, Werder M, Thuahnai ST, Han CH, Duong P, Williams DL, Wettstein P, Schulthess G, Phillips MC and Hauser H, 2005. Class B scavenger receptor-mediated intestinal absorption of dietary beta-carotene and cholesterol. *Biochemistry*, 44, 4517-4525.
- van Bennekum AM, Kako Y, Weinstock PH, Harrison EH, Deckelbaum RJ, Goldberg IJ and Blaner WS, 1999. Lipoprotein lipase expression level influences tissue clearance of chylomicron retinyl ester. *Journal of Lipid Research*, 40, 565-574.
- van Bennekum AM, Fisher EA, Blaner WS and Harrison EH, 2000. Hydrolysis of retinyl esters by pancreatic triglyceride lipase. *Biochemistry*, 39, 4900-4906.
- van Bennekum AM, Wei S, Gamble MV, Vogel S, Piantedosi R, Gottesman M, Episkopou V and Blaner WS, 2001. Biochemical basis for depressed serum retinol levels in transthyretin-deficient mice. *Journal of Biological Chemistry*, 276, 1107-1113.
- van Buuren S, Schönbeck Y and van Dommelen P, 2012. Collection, collation and analysis of data in relation to reference heights and reference weights for female and male children and adolescents (0-18 years) in the EU, as well as in relation to the age of onset of puberty and the age at which different stages of puberty are reached in adolescents in the EU. EFSA supporting publication 2012:EN-255, 59 pp.
- van Heek M, Farley C, Compton DS, Hoos L and Davis HR, 2001. Ezetimibe selectively inhibits intestinal cholesterol absorption in rodents in the presence and absence of exocrine pancreatic function. *British Journal of Pharmacology*, 134, 409-417.
- van het Hof K, Brouwer I, West C, Haddeman E, Steegers-Theunissen R, van Dusseldorp M, Weststrate J, Eskes T and Hautvast J, 1999. Bioavailability of lutein from vegetables is 5 times higher than that of beta-carotene. *American Journal of Clinical Nutrition*, 70, 261-268.
- van Lieshout M, West CE, Muhilal, Permaesih D, Wang Y, Xu X, van Breemen RB, Creemers AF, Verhoeven MA and Lugtenburg J, 2001. Bioefficacy of beta-carotene dissolved in oil studied in children in Indonesia. *American Journal of Clinical Nutrition*, 73, 949-958.
- Van Loo-Bouwman CA, Naber TH and Schaafsma G, 2014. A review of vitamin A equivalency of beta-carotene in various food matrices for human consumption. *British Journal of Nutrition*, 111, 2153-2166.
- van Rossum CTM, Fransen HP, Verkaik-Kloosterman J, Buurma-Rethans EJM and Ocké MC, 2011. Dutch National Food Consumption Survey 2007-2010: Diet of children and adults aged 7 to 69 years. RIVM Report number: 350050006/2011, National Institute for Public Health and the Environment, 143 pp.

- van Vliet T, van Schaik F, Schreurs WH and van den Berg H, 1996. In vitro measurement of beta-carotene cleavage activity: methodological considerations and the effect of other carotenoids on beta-carotene cleavage. *International Journal for Vitamin and Nutrition Research*, 66, 77-85.
- von Lintig J, 2010. Colors with functions: elucidating the biochemical and molecular basis of carotenoid metabolism. *Annual Review of Nutrition*, 30, 35-56.
- von Lintig J, Kiser PD, Golczak M and Palczewski K, 2010. The biochemical and structural basis for trans-to-cis isomerization of retinoids in the chemistry of vision. *Trends in Biochemical Sciences*, 35, 400-410.
- von Lintig J, 2012. Provitamin A metabolism and functions in mammalian biology. *American Journal of Clinical Nutrition*, 96, 1234S-1244S.
- von Reinersdorff D, Green MH and Green JB, 1998. Development of a compartmental model describing the dynamics of vitamin A metabolism in men. *Advances in Experimental Medicine and Biology*, 445, 207-223.
- Walczyk T, Davidsson L, Rossander-Hulthen L, Hallberg L and Hurrell RF, 2003. No enhancing effect of vitamin A on iron absorption in humans. *American Journal of Clinical Nutrition*, 77, 144-149.
- Wald G, 1968. Molecular basis of visual excitation. *Science*, 162, 230-239.
- Wang DQ, 2003. New concepts of mechanisms of intestinal cholesterol absorption. *Annals of Hepatology*, 2, 113-121.
- Wang Z, Yin S, Zhao X, Russell RM and Tang G, 2004. beta-Carotene-vitamin A equivalence in Chinese adults assessed by an isotope dilution technique. *British Journal of Nutrition*, 91, 121-131.
- Wang Z, Joshi AM, Ohnaka K, Morita M, Toyomura K, Kono S, Ueki T, Tanaka M, Kakeji Y, Maehara Y, Okamura T, Ikejiri K, Futami K, Maekawa T, Yasunami Y, Takenaka K, Ichimiya H and Terasaka R, 2012. Dietary intakes of retinol, carotenes, vitamin C, and vitamin E and colorectal cancer risk: the Fukuoka colorectal cancer study. *Nutrition and Cancer*, 64, 798-805.
- Ward BJ, 1996. Retinol (vitamin A) supplements in the elderly. *Drugs and Aging*, 9, 48-59.
- Wei S, Lai K, Patel S, Piantedosi R, Shen H, Colantuoni V, Kraemer FB and Blaner WS, 1997. Retinyl ester hydrolysis and retinol efflux from BFC-1beta adipocytes. *Journal of Biological Chemistry*, 272, 14159-14165.
- West CE, Eilander A and van Lieshout M, 2002. Consequences of revised estimates of carotenoid bioefficacy for dietary control of vitamin A deficiency in developing countries. *Journal of Nutrition*, 132, 2920S-2926S.
- White JA, Guo YD, Baetz K, Beckett-Jones B, Bonasoro J, Hsu KE, Dilworth FJ, Jones G and Petkovich M, 1996. Identification of the retinoic acid-inducible all-trans-retinoic acid 4-hydroxylase. *Journal of Biological Chemistry*, 271, 29922-29927.
- White JA, Beckett-Jones B, Guo YD, Dilworth FJ, Bonasoro J, Jones G and Petkovich M, 1997. cDNA cloning of human retinoic acid-metabolizing enzyme (hP450RAI) identifies a novel family of cytochromes P450. *Journal of Biological Chemistry*, 272, 18538-18541.
- WHO (World Health Organization), 1982. Control of vitamin A deficiency and xerophthalmia. Technical Report Series 672, 70 pp.
- WHO (World Health Organization), 1996. Indicators for assessing vitamin A deficiency and their application in monitoring and evaluating intervention programmes. 66 pp.
- WHO (World Health Organization), 2009. Global prevalence of vitamin A deficiency in populations at risk 1995-2006. WHO Global database on Vitamin A deficiency. 68 pp.
- WHO (World Health Organization), 2011. Serum retinol concentrations for determining the prevalence of vitamin A deficiency in populations. WHO/NMH/NHD/MNM/11.3, 5 pp.

- WHO (World Health Organization), 2012. Priorities in the assessment of vitamin A and iron status in populations. Panama City, Panama, 15-17 september 2010. 80 pp.
- WHO Multicentre Growth Reference Study Group (World Health Organization), 2006. WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development. 312 pp.
- WHO/FAO (World Health Organization/Food and Agriculture Organization of the United Nations), 2004. Vitamin and mineral requirements in human nutrition: report of a joint FAO/WHO expert consultation, Bangkok, Thailand, 21-30 September 1998. 341 pp.
- Wirtz GM, Bornemann C, Giger A, Müller RK, Schneider H, Schlotterbeck G, Schiefer G and Woggon W-D, 2001. The substrate specificity of β,β -carotene 15,15'-monooxygenase. *Helvetica Chimica Acta*, 84, 2301-2315.
- Wu T, Ni J and Wei J, 2005. Vitamin A for non-measles pneumonia in children. *Cochrane Database of Systematic Reviews*, 3, CD003700.
- Xi J and Yang Z, 2008. Expression of RALDHs (ALDH1As) and CYP26s in human tissues and during the neural differentiation of P19 embryonal carcinoma stem cell. *Gene Expression Patterns*, 8, 438-442.
- Xu X, Yu E, Liu L, Zhang W, Wei X, Gao X, Song N and Fu C, 2013. Dietary intake of vitamins A, C, and E and the risk of colorectal adenoma: a meta-analysis of observational studies. *European Journal of Cancer Prevention*, 22, 529-539.
- Yan W, Jang GF, Haeseleer F, Esumi N, Chang J, Kerrigan M, Campochiaro M, Campochiaro P, Palczewski K and Zack DJ, 2001. Cloning and characterization of a human beta,beta-carotene-15,15'-dioxygenase that is highly expressed in the retinal pigment epithelium. *Genomics*, 72, 193-202.
- Yong LC, Brown CC, Schatzkin A, Dresser CM, Slesinski MJ, Cox CS and Taylor PR, 1997. Intake of vitamins E, C, and A and risk of lung cancer. The NHANES I epidemiologic followup study. First National Health and Nutrition Examination Survey. *American Journal of Epidemiology*, 146, 231-243.
- Young JF, Luecke RH, Pearce BA, Lee T, Ahn H, Baek S, Moon H, Dye DW, Davis TM and Taylor SJ, 2009. Human organ/tissue growth algorithms that include obese individuals and black/white population organ weight similarities from autopsy data. *Journal of Toxicology and Environmental Health. Part A*, 72, 527-540.
- Zablotska LB, Gong Z, Wang F, Holly EA and Bracci PM, 2011. Vitamin D, calcium, and retinol intake, and pancreatic cancer in a population-based case-control study in the San Francisco Bay area. *Cancer Causes & Control*, 22, 91-100.
- Zhang M, Lee AH and Binns CW, 2004. Reproductive and dietary risk factors for epithelial ovarian cancer in China. *Gynecologic Oncology*, 92, 320-326.
- Zhang X, Dai B, Zhang B and Wang Z, 2012. Vitamin A and risk of cervical cancer: a meta-analysis. *Gynecologic Oncology*, 124, 366-373.
- Zile MH, Inhorn RC and DeLuca HF, 1982. Metabolites of all-trans-retinoic acid in bile: identification of all-trans- and 13-cis-retinoyl glucuronides. *Journal of Biological Chemistry*, 257, 3537-3543.
- Zovich DC, Orologa A, Okuno M, Kong LW, Talmage DA, Piantedosi R, Goodman DS and Blaner WS, 1992. Differentiation-dependent expression of retinoid-binding proteins in BFC-1 beta adipocytes. *Journal of Biological Chemistry*, 267, 13884-13889.

APPENDICES

Appendix A. Prospective cohort and nested case-control studies on the association between intake of “vitamin A” and retinol and risk of bone fracture

Reference	Design	Study sample	Dietary assessment	Outcomes	Daily intake of vitamin A (µg RE/day) ^(a) and retinol (µg/day)	Other factors considered in the analysis	Results
Melhus et al. (1998) ^{(b)(c)}	Nested case-control study within the Swedish Mammography Cohort	1 120 women, aged 40–76 years, in Sweden; 247 cases/873 controls	One FFQ covering the previous six months performed at baseline. No information on inclusion of supplements	Incidence of hip fracture. Hospital discharge records	<u>Mean ± SD (range) of retinol intake</u> Cases: 960 ± 480 (260–3 210) µg Controls: 880 ± 430 (260–5 510) µg	Energy intake, BMI, age at menopause, lifetime physical activity, smoking status, hormone replacement therapy, diabetes mellitus, oral contraceptive or cortisone use, previous osteoporotic fracture, intake of iron, magnesium, vitamin C, and calcium	<u>Retinol</u> Multivariate OR = 2.05 (95 % CI = 1.05–3.98) with a retinol intake >1 500 µg/day (highest category) compared with ≤ 500 µg/day (lowest category) p for trend = 0.006 <u>β-Carotene</u> No association found (data not shown)
Feskanich et al. (2002) ^{(b)(c)}	Prospective study, 18 years' follow-up (Nurses' Health Study, 1980–1998)	72 337 postmenopausal women aged 34–77 years in the USA	Semi-quantitative FFQ performed five times over study duration. Mean intake value determined from the mean of the five FFQs. Retinol and carotenoid content of foods from US Department of Agriculture and National Cancer Institute sources. Use of brand-specific supplements included	Incidence of hip fracture. Self-reported by questionnaire every two years	<u>Quintiles of vitamin A</u> <i>From food only</i> (n = 34 386, excluding supplement users) Q1: < 1 000; Q2: 1 000–1 299; Q3: 1 300–1 599; Q4: 1 600–1 999; Q5: ≥ 2 000 µg RE <i>From food and supplements</i> (n = 72 337) Q1: < 1 250; Q2: 1 250–1 699; Q3: 1 700–2 249; Q4: 2 250–2 999; Q5: ≥ 3 000 µg RE <u>Quintiles of retinol intake</u> <i>From food only</i> (n = 34 386, excluding supplement users) Q1: < 400; Q2: 400–549; Q3:	Age, follow-up cycle, intake of calcium, vitamin D, vitamin K, protein, alcohol and caffeine, smoking status, number of cigarettes smoked per day, use of postmenopausal hormones, body weight, hours of physical activity a day, use of thiazide diuretics	<u>Vitamin A</u> <i>Food only (excluding supplement users)</i> No association (multivariate RR). <i>Food and supplements</i> Multivariate RR = 1.48 (95 % CI = 1.05–2.07) with a vitamin A intake ≥ 3 000 µg RE/day (Q5) compared with < 1 250 µg RE/day (Q1) p for trend = 0.003 <u>Retinol</u> <i>Food only (excluding supplement users)</i> Multivariate RR = 1.69 (95 % CI = 1.05–2.74) with a retinol intake ≥ 1 000 µg/day (Q5) compared with

Reference	Design	Study sample	Dietary assessment	Outcomes	Daily intake of vitamin A ($\mu\text{g RE/day}$) ^(a) and retinol ($\mu\text{g/day}$)	Other factors considered in the analysis	Results
					550–699; Q4: 700–999; Q5: $\geq 1\,000\ \mu\text{g}$ <i>From food and supplements</i> ($n = 72\,337$) Q1: < 500; Q2: 500–849; Q3: 850–1 299; Q4: 1 300–1 999; Q5: $\geq 2\,000\ \mu\text{g}$		< 400 $\mu\text{g/day}$ (Q1) p for trend = 0.05 <i>Food and supplements</i> Multivariate RR = 1.89 (95 % CI = 1.33–2.68) with a retinol intake $\geq 2\,000\ \mu\text{g/day}$ (Q5) compared to < 500 $\mu\text{g/day}$ (Q1) Multivariate RR = 1.43 (95 % CI = 1.04–1.96) with a retinol intake of 1 300–1 999 $\mu\text{g/day}$ (Q4) compared with < 500 $\mu\text{g/day}$ (Q1) p for trend = < 0.001 <u>β-Carotene</u> No association (multivariate RR)
Michaelsson et al. (2003) ^(e)	Prospective study, 30 years' follow-up	1 221 men, aged 49–51 years, in Sweden	Seven-day dietary assessment, 20 years after entry into study. Food composition from Swedish National Food Administration database. Use of brand-specific supplements included	Incidence of any fracture. Hospital discharge register	Not provided	Energy intake	<u>Retinol</u> <i>Food only</i> Rate ratio (energy-adjusted) = 2.00 (95 % CI = 1.00–3.99) for any fracture with retinol intake > 1 500 $\mu\text{g/day}$ (Q5) compared to < 530 $\mu\text{g/day}$ (Q1). <i>Food and supplements</i> Rate ratio (energy-adjusted) = 1.99 (95 % CI = 0.98–4.01) for any fracture for Q5 (no value reported) vs. Q1 (no value reported)
Lim et al. (2004) ^(e)	Prospective study, 9.5 years' follow-up (Iowa Women's Health Study 1986–1997)	34 703 postmenopausal women, aged 55–69 years, in the USA	One semi-quantitative FFQ performed at baseline. Use of brand-specific supplements included	Incidence of hip and non-hip fracture. Self-reported by questionnaire at the end of the follow-up period	<u>Mean (range) for each quintile of vitamin A intake (in IU)</u> <i>From food only (n = 22 410, excluding supplement users)</i> Q1: 4 440 (221–5 975); Q2: 7 223 (5 976–8 544); Q3: 10 043 (8 545–11 699); Q4:	<u>For hip fracture</u> Age, BMI, waist-to-hip ratio, diabetes mellitus, physical activity, occurrence of past irregular menstrual duration, steroid medication, oestrogen	<u>Vitamin A and retinol</u> No association (multivariate RR) between vitamin A or retinol intake, from supplements only, food and supplements, or food only (excluding supplement users), and risk of hip fracture or risk of all

Reference	Design	Study sample	Dietary assessment	Outcomes	Daily intake of vitamin A (µg RE/day) ^(a) and retinol (µg/day)	Other factors considered in the analysis	Results
					<p>13 793 (11 700–16 431); Q5: 24 163 (16 432–215 392) IU <i>From food and supplements (n = 34 703)</i></p> <p>Q1: 5 113 (221–7 055); Q2: 8 771 (7 056–10 484); Q3: 12 256 (10 485–14 209); Q4: 16 764 (14 210–19 892); Q5: 29 239 (19 893–236 991) IU</p> <p><u>Mean (range) for each quintile of retinol intake</u> <i>From food only (n = 22 410, excluding supplement users)</i> Q1: 744 (28–1 086); Q2: 1 424 (1 087–1 790); Q3: 2 356 (1 791–3 260); Q4: 3 967 (3 261–4 660); Q5: 6 878 (4 661–209 573) IU <i>From food and supplements (n = 34 703)</i> Q1: 912 (28–1405); Q2: 2 031 (1 406–2 952); Q3: 3 857 (2 953–4 655); Q4: 5 768 (4 656–7 001); Q5: 12 610 (7 002–211 051) IU</p>	<p>replacement, energy intake</p> <p><u>For all fractures</u> Age, BMI, waist-to-hip ratio, diabetes mellitus, cirrhosis, past irregular menstrual duration, thyrotropic, sedative, antiepileptic or diuretic medications, education, alcohol use and energy intake</p>	fractures
Rejnmark et al. (2004)	Nested case–control study	1 141 perimenopausal women aged 45–58 years in Denmark; 163 cases/978 controls	Four- or seven-day food record at baseline and after five years. Composition data from official Danish food tables. Use of supplements included	Incidence of fractures. Self-reported, confirmed by hospital discharge records	<p><u>Median (interquartile range) of vitamin A intake</u> <i>From food only</i> Cases: 1 150 (730–1 720) µg RE Controls: 1 140 (800–1 660) µg RE <i>From food and supplements</i> Cases: 1 730 (1 280–2 380) µg RE Controls: 1 710 (1 290–2 260) µg RE</p>		<p><u>Vitamin A, retinol and β-carotene</u> No association (multivariate OR) between vitamin A, retinol or β-carotene intake, from food only or food and supplements, and risk of fracture</p>

Reference	Design	Study sample	Dietary assessment	Outcomes	Daily intake of vitamin A ($\mu\text{g RE/day}$) ^(a) and retinol ($\mu\text{g/day}$)	Other factors considered in the analysis	Results
					<p><u>Median (interquartile range) of retinol intake</u> <i>From food only</i> Cases: 510 (350–700) μg Controls: 520 (380–740) μg <i>From food and supplements</i> Cases: 1 190 (700–1 420) μg Controls: 1 210 (740–1 430) μg</p>		
Caire-Juvera et al. (2009)	Prospective study. 6.6 years' follow-up (Women's Health Initiative Observational Study 1993–2005)	75 747 postmenopausal women, mean age at baseline 63.6 years, in the USA	FFQ at baseline and at three-year follow-up. Mean intake value determined from the mean of the two FFQs. Retinol and carotenoid content of foods from the University of Minnesota Nutrition Coding Center nutrient database. Use of brand-specific supplements included	Incidence of hip and non-hip fracture. Questionnaire every year from participants or proxy respondents. Hip fractures were confirmed by medical records	<p><u>Quintiles of vitamin A</u> <i>From food and supplements</i> Q1: < 5 055; Q2: 5 055–5 824; Q3: 5 825–6 550; Q4: 6 551–7 507; Q5: \geq 7 508 $\mu\text{g RE}$</p> <p><u>Mean \pm SD intake of retinol, for each quintile of vitamin A intake</u> <i>From food and supplements</i> Q1: 412 \pm 187; Q2: 727 \pm 284; Q3: 983 \pm 341; Q4: 1 227 \pm 407; Q5: 1 968 \pm 1 266 μg</p>	Age, intake of protein, vitamin D, vitamin K, calcium, caffeine, and alcohol, BMI, hormone therapy, smoking, metabolic equivalents hours per week, ethnicity and region of clinical centre	<p><u>Vitamin A and retinol</u> No association (multivariate HR, including vitamin D and calcium) between vitamin A intake or retinol, from food and supplements, and risk of hip fracture or risk of total fracture</p> <p>Among the women with lower vitamin D intake (\leq 11 $\mu\text{g/day}$), there was a higher risk of total fractures in Q5 of vitamin A intake (mean 8 902 $\mu\text{g RE/day}$) compared with Q1 (mean 4 445 $\mu\text{g RE/day}$) (HR: 1.19; 95 % CI = 1.04–1.37; p for trend = 0.022) and in Q5 of retinol intake (mean 2 488 $\mu\text{g/day}$) compared with Q1 (mean 348 $\mu\text{g/day}$) (HR: 1.15; 95 % CI = 1.03–1.29; p for trend = 0.056). Given the smaller number of hip fractures, stratified analysis by vitamin D and calcium intake was not conducted</p>
Ambrosini et al. (2013)	Retrospective analysis of the Vitamin A Program	664 women and 1 658 men in Australia (99 % participants on the	Background dietary intake not assessed. Supplementation with 7 500 $\mu\text{g/day}$ of retinol	Database on hospital admissions for fracture and self-reported by	Background dietary intake not reported. Cumulative dose of retinol supplements was estimated by summing	Age, sex, smoking, BMI, medication use and previous fractures	<p><u>Retinol</u> No associations (multivariate OR) between cumulative dose of retinol and risk for any</p>

Reference	Design	Study sample	Dietary assessment	Outcomes	Daily intake of vitamin A ($\mu\text{g RE/day}$) ^(a) and retinol ($\mu\text{g/day}$)	Other factors considered in the analysis	Results
		Vitamin A Program), mean age at enrolment 55 years	as retinyl palmitate for 1–16 years (median 7 years)	questionnaire sent to all surviving participants after the end of the intervention. Self-reported fractures occurring at the spine, hip, femur, arm, ribs or wrist were classified as osteoporotic fractures	the number of days the supplement was taken between each annual follow-up, multiplying by the dose administered and adding to the previous year's total. Cumulative doses of retinol were analysed in units of 10 g. The maximum cumulative dose of retinol was 42 g, equivalent to taking 7 500 $\mu\text{g/day}$ for 15.3 years		fracture or osteoporotic fracture

BMI, body mass index; CI, confidence interval; FFQ, food frequency questionnaire; HR, hazard ratio; OR, odds ratio; Q, quintile; RR, relative risk; IU, International Unit.

(a): Unless stated otherwise.

(b): Study considered in SCF (2002).

(c): Study considered in SACN (2005).

Appendix B. Intervention and prospective cohort studies on the association between intake of “vitamin A” and retinol and measures of bone mineral content (BMC), bone mineral density (BMD) or serum markers of bone turnover

Reference	Design	Study sample	Dietary assessment	Outcomes	Daily intake of vitamin A (µg RE/day) ^(a) and retinol (µg/day)	Other factors considered in the analysis	Results
Measures of BMC, BMD							
Freudenheim et al. (1986) ^{(b)(c)}	Prospective study, four years' follow-up within a calcium supplementation trial	99 pre- and postmenopausal women, aged 35–65 years, in the USA	Seventy-two 24-hour dietary records collected for each participant over three years, including supplements. Subjects were assigned to a 500 mg calcium-supplemented or a placebo group	BMC of left arm bones (radius, humerus and ulna). By SPA. Eleven measurements, every three months for the first year and then every six months	<u>Mean ± SD (range) “vitamin A”^(d) intake from food and supplements (in IU)</u> <i>Postmenopausal calcium unsupplemented (n = 33)</i> 8 624 ± 3 553 (3 615–17 763) IU <i>Calcium supplemented (n = 34)</i> 7 619 ± 2 729 (3 256–14 624) IU	None.	<u>“Vitamin A”^(d)</u> In the postmenopausal calcium-supplemented group, there was a negative correlation between “vitamin A” intake and rate of change in ulna BMC – the correlation was not significant when one subject with very high supplemental “vitamin A” intake was omitted. No correlation was observed in the postmenopausal calcium-unsupplemented group There were no correlations in the groups of calcium-supplemented (n = 8) and calcium-unsupplemented (n = 9) premenopausal women
Houtkooper et al. (1995) ^{(b)(c)}	One year follow-up within a physical exercise trial	66 premenopausal women, aged 28–39 years, in the USA	Dietary records over 4–12 randomly assigned days. Vitamin supplements not included. All subjects were administered a 500-mg calcium supplement	BMD of total body, lumbar vertebrae 2-4, femoral neck, Ward's triangle, trochanter. By DXA. Four measurements, at baseline and months 5, 12 and 18	<u>Mean ± SD “vitamin A”^(d)</u> <i>From food only</i> 1 220 ± 472 µg RE	Fat mass at baseline and change in fat mass over one year, exercise status.	<u>“Vitamin A”^(d)</u> Significant variables in models predicting total body BMD slope included initial fat mass and fat mass slope plus either “vitamin A” ^(d) intake (R ² = 0.31) or β-carotene intake (R ² = 0.28)
Promislow et al. (2002) ^(c)	Prospective study, with four years' follow-up, within the Ranchi	570 women and 388 men, aged 55–92 years at baseline, in the USA	FFQ at baseline. Supplement use included	BMD of total hip, femoral neck, lumbar spine. By DXA. Two	<u>Mean ± SD of retinol intake</u> <i>From food only</i> Women: 497 ± 460 µg	Age, weight change, BMI, calcium intake, diabetes status,	<u>Retinol</u> No association between retinol intake and BMD at baseline or BMD change when supplement

Reference	Design	Study sample	Dietary assessment	Outcomes	Daily intake of vitamin A (µg RE/day) ^(a) and retinol (µg/day)	Other factors considered in the analysis	Results
	Bernardo Heart and Chronic Disease Study			measurements, taken at baseline and follow-up	Men: 624 ± 585 µg <i>From food and supplements</i> Women: 1 247 ± 1 573 µg Men: 1 242 ± 1 442 µg	menopausal status, exercise, smoking status, alcohol use, thiazide drug use, thyroid hormone use, steroid use, oestrogen use, supplemental retinol	users and non-users were pooled <i>For supplement users only</i> Women: a significant negative association was found between retinol intake and BMD at the femoral neck (p = 0.02) and total spine (p = 0.03) measured at follow-up and for BMD change at femoral neck (p = 0.05) and total hip (p = 0.02) Men: no significant association
Macdonald et al. (2004)	Prospective study, within the Aberdeen Prospective Osteoporosis Screening Study, 5–7 years' follow-up	891 women, aged 45–55 years at baseline, in the UK	FFQ at baseline and five years later. Composition data from McCance and Widdowson's food composition tables. Use of brand-specific supplements included	BMD of lumbar spine, femoral neck By DXA Two measurements, at baseline and follow-up	<u>Vitamin A intake</u> Not reported. <u>Mean ± SD (range, median) of retinol intake</u> <i>From food only</i> Baseline: 820 ± 602 (39–4 354, 588) µg Follow-up: 665 ± 513 (70–5 237, 480) µg <i>From food and supplements</i> Baseline: 924 ± 666 (85–4 354, 702) µg Follow-up: 882 ± 654 (70–5 237, 627) µg	Energy intake, age, weight, annual percentage change in weight, height, smoking status, socioeconomic status, physical activity level, baseline BMD measurement, menopausal status and hormone replacement therapy use	<u>Vitamin A</u> In multiple regression analysis, vitamin A intake from food only was a weak but significant negative predictor of femoral neck BMD change (variation explained: 0.3 %, coefficient (95 % CI): –1.24 (–2.47 to 0.17), p = 0.047). No significant relation when intake from supplements was included <u>Retinol</u> In multiple regression analysis, retinol intake from food only was a weak but significant negative predictor of femoral neck BMD change (variation explained: 0.4 %, coefficient (95 % CI): –1.73 (–3.20 to –0.30), p = 0.018). No significant relation when intake from supplements was included
Rejnmark et al. (2004)	Prospective study, five years' follow-up within the Danish Osteoporosis Prevention Study	1 694 perimenopausal women, aged 45–58 years, in Denmark	Four- or seven-day food record at baseline and after five years. Intake at baseline was	BMD of lumbar spine, femoral neck. By DXA. Two measurements, at baseline and five-	<u>Median (interquartile range) vitamin A intake (baseline)</u> <i>From food only</i> 1 150 (800–1 730) µg RE <i>From food and</i>	Age, years postmenopausal, hormone therapy, previous fracture, body weight, baseline BMD,	<u>Vitamin A and retinol</u> Multiple regression analysis showed no association between baseline vitamin A or retinol intake, from food only or food and supplements, and change in BMD

Reference	Design	Study sample	Dietary assessment	Outcomes	Daily intake of vitamin A ($\mu\text{g RE/day}$) ^(a) and retinol ($\mu\text{g/day}$)	Other factors considered in the analysis	Results
			considered in the analysis. Composition data from official Danish food tables. Use of supplements included	year follow-up	<i>supplements</i> 1 740 (1 290–2 360) $\mu\text{g RE}$ <u>Median (interquartile range) retinol intake (baseline)</u> <i>From food only</i> 530 (390–750) μg <i>From food and supplements</i> 1 210 (680–1 450) μg	physical activity, energy intake, intake of calcium, vitamin D, alcohol, smoking status, use of thiazide or loop diuretics, thyroid hormones, antipsychotic/ anxiolytic/ antidepressant drugs, diagnosis of thyrotoxicosis, diabetes mellitus	at any site <u>β-Carotene</u> No association between β -carotene intake, from food only, and change in BMD at any site
Measures of serum markers of bone turnover							
Kawahara et al. (2002) ^{(b) (c)}	Randomised single-blind trial lasting six weeks	80 men, aged 18–58 years, in the USA	Subjects were assigned to 7 576 μg retinyl palmitate/day or placebo. Background retinol intake not assessed	Serum osteocalcin, bone-specific alkaline phosphatase, N-telopeptide of type 1 collagen/ Blood sampled at baseline and weeks 2, 4 and 6	Not reported.		<u>Retinol</u> Supplementation did not affect serum osteocalcin, bone-specific alkaline phosphatase or N-telopeptide of type 1 collagen

BMC, bone mineral content; BMD, bone mineral density; BMI, body mass index; CI, confidence interval; DXA, dual energy X-ray absorptiometry; FFQ, food frequency questionnaire; IU, International Unit; SPA, single-photon absorptiometry.

(a): Unless stated otherwise.

(b): Study considered in SCF (2002).

(c): Study considered in SACN (2005).

(d): It is unclear from the article whether it refers to vitamin A or retinol only.

Appendix C. Retinol concentration in breast milk from mothers of term infants

Reference	Number of women	Country	Maternal vitamin A intake	Stage of lactation (time post partum)	Concentration ($\mu\text{g/L}$)			Methods ^(a)
					Mean \pm SD	Median	Range	
Canfield et al. (2003)	53	Australia	Not reported. Mothers who were taking supplements containing carotenoids or vitamin A (> 8 000 IU/day) were excluded	Mature milk (months 2–12)	311 \pm 16 (SE)			Single complete breast expression by electric breast pump collected mid-afternoon from each mother. Samples were collected from the breast from which the infant had most recently fed. Samples were saponified before analysis of retinol by HPLC
	55	Canada		Mature milk (months 2–12)	340 \pm 19 (SE)			
	50	UK		Mature milk (months 2–12)	301 \pm 14 (SE)			
	49	USA		Mature milk (months 2–12)	352 \pm 25 (SE)			
Schweigert et al. (2004)	21	Germany	Not reported. Mothers taking supplements containing carotenoids or vitamin A were excluded	Colostrum (days 4 \pm 2)	1 532 \pm 725			Total milk volume of one breast was collected. Samples were saponified before analysis of retinol by HPLC
	21			Mature milk (days 19 \pm 2)	831 \pm 321			
Schulz et al. (2007)	26	Germany	Mean \pm SD: Retinol intake: 0.95 \pm 0.64 mg/day Carotenoid intake: 6.9 \pm 3.6 mg/day Total vitamin A intake: 2.11 \pm 0.89 mg RE/day. By FFQ. Mothers taking prenatally supplements with > 2 000 IU/day of vitamin A or > 2 mg/day of β -carotene were excluded	Colostrum (days 1–2)	1 106 \pm 851			Samples collected by hand expression or electric pump up to a volume of 4 mL, collected on one or more occasions. Samples were saponified before analysis of retinol by HPLC
Tokusoglu et al. (2008)	92	Turkey	Not reported	Mature milk (days 60–90)	815 \pm 120.6			Milk samples (10 mL) collected from both breasts by hand expression, at least two hours after previous breastfeeding. Samples were saponified before analysis of retinol by HPLC

Reference	Number of women	Country	Maternal vitamin A intake	Stage of lactation (time post partum)	Concentration (µg/L)			Methods ^(a)
					Mean ± SD	Median	Range	
Duda et al. (2009) ^(b)	30	Poland	Mean ± SD: “vitamin A-equivalent” intake: 1 012 ± 735 µg/day β-carotene intake: 2 096 ± 2 465 µg/day By 24-hour recall (repeated on three consecutive days)	Mature milk (months 2–4)	571 ± 500	294	157–1 424	Milk samples expressed by hand or using a sterile pump one or two hours prior to actual feeding of the baby. Samples were saponified before analysis of retinol by HPLC
Orhon et al. (2009)	20	Turkey	Mean ± SEM: 4 965.2 ± 538.5 IU/day By 5-day dietary record Significant correlation between dietary vitamin A intake and retinol concentration of breast milk (r = 0.621, p = 0.006). No correlation between dietary vitamin A intake and β-carotene concentration in breast milk	Transitional milk (day 7)	2 463 ± 200 (SE)			Milk samples (5 mL) were collected from each breast using an electric pump. Treatment of the samples not described. Retinol analysed by HPLC
Kasparova et al. (2012) ^(b)	12	Czech Republic	Not reported	Mature milk: months 1–2	458 ± 286			Milk samples obtained from a university hospital; method of expression not described. Samples were saponified before analysis of retinol by HPLC
				Mature milk: months 3–4	315 ± 258			
				Mature milk: months 5–6	229 ± 115			
				Mature milk: months 9–12	172 ± 115			

Studies were identified by a comprehensive literature search for publications from January 2000 to January 2014 (LASER Analytica, 2014). Note that if studies did not report whether infants were born at term or not it was presumed that infants were born at term.

HPLC, high-performance liquid chromatography; IU, International Unit.

(a): Determination of total breast milk retinol requires saponification (typically with alcoholic potassium hydroxide (KOH)) and retinol is then extracted with an organic solvent, usually hexanes, before HPLC analysis (Tanumihardjo and Penniston, 2002).

(b): It was not reported whether the infants were born at term or not.

Appendix D. Dietary surveys in the updated EFSA Comprehensive European Food Consumption Database included in the nutrient intake calculation and number of subjects in the different age classes

Country	Dietary survey	Year	Method	Days	Number of subjects ^(a)					
					Children 1 to < 3 years	Children 3 to < 10 years	Children 10 to < 18 years	Adults 18 to < 65 years	Adults 65 to < 75 years	Adults ≥ 75 years
Finland/1	DIPP	2000–2010	Dietary record	3	500	750				
Finland/2	NWSSP	2007–2008	48-hour dietary recall ^(b)	2 × 2 ^(b)			306			
Finland/3	FINDIET2012	2012	48-hour dietary recall ^(b)	2 ^(b)				1 295	413	
France	INCA2	2006–2007	Dietary record	7		482	973	2 276	264	84
Germany/1	EsKiMo	2006	Dietary record	3		835	393			
Germany/2	VELS	2001–2002	Dietary record	6	347	299				
Ireland	NANS	2008–2010	Dietary record	4				1 274	149	77
Italy	INRAN-SCAI 2005-06	2005–2006	Dietary record	3	36 ^(a)	193	247	2 313	290	228
Latvia	FC_PREGNANTWOMEN 2011	2011	24-hour dietary recall	2			12 ^(a)	991 ^(c)		
Netherlands	DNFCS 2007–2010	2007–2010	24-hour dietary recall	2		447	1 142	2 057	173	
Sweden	RISKMATEN	2010–2011	Dietary record (Web)	4				1 430	295	72
United Kingdom	NDNS - Rolling Programme (1–3 years)	2008–2011	Dietary record	4	185	651	666	1 266	166	139

DIPP, type 1 Diabetes Prediction and Prevention survey; DNFCS, Dutch National Food Consumption Survey; EsKiMo, Ernährungstudie als KIGGS-Modul; FINDIET, the national dietary survey of Finland; INCA, étude Individuelle Nationale de Consommations Alimentaires; INRAN-SCAI, Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione—Studio sui Consumi Alimentari in Italia; FC_PREGNANTWOMEN, food consumption of pregnant women in Latvia; NANS, National Adult Nutrition Survey; NDNS, National Diet and Nutrition Survey; NWSSP, Nutrition and Wellbeing of Secondary School Pupils; VELS, Verzehrsstudie zur Ermittlung der Lebensmittelaufnahme von Säuglingen und Kleinkindern für die Abschätzung eines akuten Toxizitätsrisikos durch Rückstände von Pflanzenschutzmitteln.

(a): 5th or 95th percentile intakes calculated from fewer than 60 subjects require cautious interpretation as the results may not be statistically robust (EFSA, 2011b) and, therefore, for these dietary surveys/age classes, the 5th and 95th percentile estimates will not be presented in the intake results.

(b): A 48-hour dietary recall comprises two consecutive days.

(c): One subject was excluded from the dataset because only one 24-hour dietary recall day was available, i.e. final n = 990.

Appendix E. Vitamin A intake in males in different surveys according to age classes and country ($\mu\text{g RE/day}$)

Age class	Country	Survey	n	Average	P5	P50	P95
1 to < 3 years	Finland	DIPP_2001_2009	245	491	116	419	1 134
	Germany	VELS	174	651	264	582	1 294
	Italy	INRAN_SCAI_2005_06	20	554	^(a) 499	^(a)	
	United Kingdom	NDNS-RollingProgrammeYears1-3	107	576	260	496	1 032
3 to < 10 years	Finland	DIPP_2001_2009	381	751	243	550	2 022
	France	INCA2	239	702	240	579	1 353
	Germany	EsKiMo	426	889	329	754	1 951
	Germany	VELS	146	685	331	656	1 271
	Italy	INRAN_SCAI_2005_06	94	873	293	618	1 475
	Netherlands	DNFCS 2007–2010	231	741	204	589	1 876
	United Kingdom	NDNS-RollingProgrammeYears1-3	326	607	245	531	1 104
10 to < 18 years	Finland	NWSSP07_08	136	776	285	644	1 391
	France	INCA2	449	758	259	635	1 475
	Germany	EsKiMo	197	949	361	803	2 213
	Italy	INRAN_SCAI_2005_06	108	891	360	688	1 766
	Netherlands	DNFCS 2007–2010	566	866	249	664	2 076
	United Kingdom	NDNS-RollingProgrammeYears1-3	340	686	236	600	1 351
18 to < 65 years	Finland	FINDIET2012	585	1 078	325	867	2 154
	France	INCA2	936	978	279	747	2 068
	Ireland	NANS_2012	634	1 023	356	891	1 864
	Italy	INRAN_SCAI_2005_06	1 068	984	345	750	1 924
	Netherlands	DNFCS 2007–2010	1 023	1 097	340	858	2 662
	Sweden	Riksmaten 2010	623	995	311	880	2 005
	United Kingdom	NDNS-RollingProgrammeYears1-3	560	930	268	768	1 847
65 to < 75 years	Finland	FINDIET2012	210	1 086	307	823	2 345
	France	INCA2	111	1 279	367	892	5 080
	Ireland	NANS_2012	72	1 243	360	1 173	2 558
	Italy	INRAN_SCAI_2005_06	133	1 036	353	772	2 058

Age class	Country	Survey	n	Average	P5	P50	P95
≥ 75 years	Netherlands	DNFCS 2007–2010	91	1 029	316	871	2 604
	Sweden	Riksmaten 2010	127	1 042	437	911	1 879
	United Kingdom	NDNS-RollingProgrammeYears1-3	75	1 423	345	1077	5 360
	France	INCA2	40	1 057	^(a)	794	^(a)
	Ireland	NANS_2012	34	992	^(a)	881	^(a)
	Italy	INRAN_SCAI_2005_06	69	949	291	722	1 635
	Sweden	Riksmaten 2010	42	1 270	^(a)	1059	^(a)
	United Kingdom	NDNS-RollingProgrammeYears1-3	56	1 353	^(a)	798	^(a)

n, number of individuals; P5, 5th percentile; P50, 50th percentile; P95, 95th percentile.

DIPP, type 1 Diabetes Prediction and Prevention survey; DNFCS, Dutch National Food Consumption Survey; EsKiMo, Ernährungstudie als KIGGS-Modul; FINDIET, the national dietary survey of Finland; INCA, étude Individuelle Nationale de Consommations Alimentaires; INRAN-SCAI, Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione—Studio sui Consumi Alimentari in Italia; NANS, National Adult Nutrition Survey; NDNS, National Diet and Nutrition Survey; NWSSP, Nutrition and Wellbeing of Secondary School Pupils; VELs, Verzehrsstudie zur Ermittlung der Lebensmittelaufnahme von Säuglingen und Kleinkindern für die Abschätzung eines akuten Toxizitätsrisikos durch Rückstände von Pflanzenschutzmitteln.

(a): 5th or 95th percentile intakes calculated from fewer than 60 subjects require cautious interpretation, as the results may not be statistically robust (EFSA, 2011b) and, therefore, for these dietary surveys/age classes, the 5th and 95th percentile estimates will not be presented in the intake results.

Appendix F. Vitamin A intake in females in different surveys according to age classes and country ($\mu\text{g RE/day}$)

Age class	Country	Survey	n	Average	P5	P50	P95
1 to < 3 years	Finland	DIPP_2001_2009	255	409	125	358	255
	Germany	VELS	174	598	240	525	174
	Italy	INRAN_SCAI_2005_06	16	446	^(a) 428	428	16
	United Kingdom	NDNS-RollingProgrammeYears1-3	78	437	182	422	78
3 to < 10 years	Finland	DIPP_2001_2009	369	647	234	501	369
	France	INCA2	243	609	230	537	243
	Germany	EsKiMo	409	793	279	715	409
	Germany	VELS	147	654	301	590	147
	Italy	INRAN_SCAI_2005_06	99	696	262	592	99
	Netherlands	DNFCS 2007–2010	216	716	203	545	216
	United Kingdom	NDNS-RollingProgrammeYears1-3	325	610	225	576	325
10 to < 18 years	Finland	NWSSP07_08	170	724	345	631	170
	France	INCA2	524	662	217	557	524
	Germany	EsKiMo	196	892	320	752	196
	Italy	INRAN_SCAI_2005_06	139	799	280	680	139
	Latvia ^(b)	FC_PREGNANTWOMEN_2011	12	1 078	^(a) 970	970	12
	Netherlands	DNFCS 2007–2010	576	713	236	573	576
	United Kingdom	NDNS-RollingProgrammeYears1-3	326	597	225	518	326
18 to < 65 years	Finland	FINDIET2012	710	960	312	799	710
	France	INCA2	1 340	979	301	713	1 340
	Ireland	NANS_2012	640	897	319	765	640
	Italy	INRAN_SCAI_2005_06	1 245	885	322	708	1 245
	Latvia ^(b)	FC_PREGNANTWOMEN_2011	990 ^(b)	1 319	375	886	990
	Netherlands	DNFCS 2007–2010	1 034	906	268	690	1 034
	Sweden	Riksmaten 2010	807	958	379	854	807
	United Kingdom	NDNS-RollingProgrammeYears1-3	706	891	266	697	706
65 to < 75 years	Finland	FINDIET2012	203	913	314	730	203
	France	INCA2	153	1 281	408	874	153

Age class	Country	Survey	n	Average	P5	P50	P95
	Ireland	NANS_2012	77	1 041	345	927	77
	Italy	INRAN_SCAI_2005_06	157	873	329	736	157
	Netherlands	DNFCS 2007–2010	82	905	317	712	82
	Sweden	Riksmaten 2010	168	1 159	373	875	168
	United Kingdom	NDNS-RollingProgramme Years 1-3	91	1 139	354	839	91
≥ 75 years	France	INCA2	44	1 498	^(a)	740	44
	Ireland	NANS_2012	43	1 050	^(a)	922	43
	Italy	INRAN_SCAI_2005_06	159	816	308	706	159
	Sweden	Riksmaten 2010	30	1 331	^(a)	987	30
	United Kingdom	NDNS-RollingProgramme Years 1-3	83	991	374	771	83

n, number of individuals; P5, 5th percentile; P50, 50th percentile; P95, 95th percentile.

DIPP, type 1 Diabetes Prediction and Prevention survey; DNFCS, Dutch National Food Consumption Survey; EsKiMo, Ernährungstudie als KIGGS-Modul; FINDIET, the national dietary survey of Finland; INCA, étude Individuelle Nationale de Consommations Alimentaires; INRAN-SCAI, Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione - Studio sui Consumi Alimentari in Italia; FC_PREGNANTWOMEN, food consumption of pregnant women in Latvia; NANS, National Adult Nutrition Survey; NDNS, National Diet and Nutrition Survey; NWSSP, Nutrition and Wellbeing of Secondary School Pupils; VELs, Verzehrsstudie zur Ermittlung der Lebensmittelaufnahme von Säuglingen und Kleinkindern für die Abschätzung eines akuten Toxizitätsrisikos durch Rückstände von Pflanzenschutzmitteln.

(a): 5th or 95th percentile intakes calculated from fewer than 60 subjects require cautious interpretation, as the results may not be statistically robust (EFSA, 2011b) and, therefore, for these dietary surveys/age classes, the 5th and 95th percentile estimates will not be presented in the intake results.

(b): Pregnant women only.

Appendix G. Minimum and maximum % contribution of different food groups to vitamin A intake in males

Food groups	Age					
	1 to < 3 years	3 to < 10 years	10 to < 18 years	18 to < 65 years	65 to < 75 years	≥ 75 years
Additives, flavours, baking and processing aids	0	0	0	0	0	0
Alcoholic beverages	0	0	< 0.1	< 0.1	< 0.1	< 0.1
Animal and vegetable fats and oils	2.2–9.9	3.2–18.2	4.4–27.3	3.8–21.9	3.1–22.7	3.2–20.3
Coffee, cocoa, tea and infusions	0–0.1	< 0.1–0.3	< 0.1–0.4	< 0.1–1.6	< 0.1–1.6	0–1.2
Composite dishes	0.5–11.4	0.6–11.8	0.8–14	0.4–24.3	0.5–19.3	0.3–19.4
Eggs and egg products	1.2–2.8	1–6.6	0.9–6.3	0.9–4.6	0.6–4.3	1–4.2
Fish, seafood, amphibians, reptiles and invertebrates	0.1–0.4	0.1–1	0.1–1.1	0.2–1.5	0.6–1.7	0.5–1.4
Food products for young population	4.9–10.2	< 0.1–1.4	< 0.1	< 0.1	–	–
Fruit and fruit products	0.9–8.9	0.5–3.2	0.4–2.2	0.3–3.4	0.5–4.2	0.4–4
Fruit and vegetable juices and nectars	0.2–9.4	1–10.4	1.1–9.1	0.6–5.4	0.3–2.9	0.1–3.2
Grains and grain-based products	0.3–7.2	0.1–9	0.2–10	3–6.5	2.7–6.1	2.9–6.2
Human milk	< 0.1–3.8	–	–	–	–	–
Legumes, nuts, oilseeds and spices	0.3–1	0.1–0.7	0.1–0.8	0.2–1.3	0.3–0.6	0.4–0.8
Meat and meat products	0.7–10	5.1–24.5	8.4–16.6	7.4–25.1	14.6–32.5	3.4–38.4
Milk and dairy products	11.6–31.8	14.7–24.1	16.9–23.8	14–18.3	10.7–16.5	11.7–17.8
Products for non-standard diets, food imitates and food supplements or fortifying agents	0–0.1	0–0.1	< 0.1–0.2	< 0.1–0.4	< 0.1–0.5	0
Seasoning, sauces and condiments	< 0.1–2.1	< 0.1–6.2	< 0.1–5.7	< 0.1–5.4	< 0.1–3.6	< 0.1–2.6
Starchy roots or tubers and products thereof, sugar plants	< 0.1–0.3	< 0.1–0.9	< 0.1–0.8	< 0.1–0.9	< 0.1–1.4	< 0.1–0.3
Sugar, confectionery and water-based sweet desserts	< 0.1–0.5	0.1–1.1	0.1–1.1	< 0.1–0.5	< 0.1–0.2	< 0.1–0.1
Vegetables and vegetable products	23.6–58.2	25.3–38.2	19–44.5	15.3–48.5	16.8–52.2	20–49.5
Water and water-based beverages	0	< 0.1–0.1	< 0.1–0.1	< 0.1–0.1	0	< 0.1–0.1

“–” means that there was no consumption event of the food group in the age and sex group considered, whereas “0” means that there were some consumption events, but that the food group does not contribute to the intake of the nutrient considered, for the age and sex group considered.

Appendix H. Minimum and maximum % contribution of different food groups to vitamin A intake in females

Food groups	Age					
	1 to < 3 years	3 to < 10 years	10 to < 18 years	18 to < 65 years	65 to < 75 years	≥ 75 years
Additives, flavours, baking and processing aids	0	0	0	0	0	0
Alcoholic beverages	0	0	0	< 0.1–0.2	< 0.1–0.1	0–0.3
Animal and vegetable fats and oils	2–11.6	3.9–18.4	3.7–25	3.6–18.6	3.3–17.7	3.1–15.9
Coffee, cocoa, tea and infusions	0–0.1	< 0.1–0.2	< 0.1–0.4	< 0.1–1.4	< 0.1–1.5	< 0.1–0.9
Composite dishes	0.1–12.8	0.7–11.4	0.4–15.6	0.4–24.7	0.4–16.7	0.3–19.4
Eggs and egg products	0.8–3.4	1–6.5	0.8–6.4	1–4	0.8–3.8	0.6–4.4
Fish, seafood, amphibians, reptiles and invertebrates	0.1–0.6	< 0.1–0.7	0.2–1.4	0.2–1.2	0.2–1.2	0.4–0.7
Food products for young population	4–16.2	< 0.1–0.6	< 0.1–0.1	< 0.1–0.1	–	0.1
Fruit and fruit products	1–9.2	0.6–2.9	0.5–4.8	0.4–4.5	0.7–5.4	0.6–5.8
Fruit and vegetable juices and nectars	0.2–8.4	0.9–8.6	1.3–10.9	0.7–4.2	0.7–3.8	0.2–4.9
Grains and grain-based products	0.4–6.5	0.1–9.1	0.1–9.8	2.7–6	2.9–4.5	2.9–4.4
Human milk	< 0.1	–	–	–	–	–
Legumes, nuts, oilseeds and spices	0.3–0.8	0.1–1	0.2–0.7	0.2–1	0.1–1	0.3–0.6
Meat and meat products	0.5–5.7	0.9–23.1	4.6–16.1	7.8–29.6	5.7–35.1	4.2–45.8
Milk and dairy products	13.4–31.3	15.7–25.9	15.9–25.4	11.7–18	8.2–16.8	9.1–18.3
Products for non-standard diets, food imitates and food supplements or fortifying agents	0–0.3	0–0.1	0–0.3	< 0.1–0.5	0–0.3	0–0.6
Seasoning, sauces and condiments	< 0.1–2.9	< 0.1–6.4	< 0.1–6.1	< 0.1–4.3	< 0.1–2.8	< 0.1–2.5
Starchy roots or tubers and products thereof, sugar plants	< 0.1–1.1	< 0.1–0.8	< 0.1–0.9	0.1–0.8	< 0.1–0.7	< 0.1–0.2
Sugar, confectionery and water-based sweet desserts	< 0.1–0.5	0.2–1.1	< 0.1–1.1	< 0.1–0.5	< 0.1–0.2	< 0.1–0.2
Vegetables and vegetable products	27.2–59.2	19.6–41.1	21–40.9	22.1–51.7	21.7–55.1	23.8–56.9
Water and water-based beverages	0	< 0.1–0.1	0–0.1	< 0.1–0.1	0	< 0.1

“–” means that there was no consumption event of the food group in the age and sex group considered, whereas “0” means that there were some consumption events, but that the food group does not contribute to the intake of the nutrient considered, for the age and sex group considered.

ABBREVIATIONS

Afssa	Agence française de sécurité sanitaire des aliments
AI	Adequate Intake
AR	Average Requirement
BCMO1	β,β -carotene-15,15'-monooxygenase 1
BMD	bone mineral density
CI	confidence interval
COMA	Committee on Medical Aspects of Food Policy
CRABP	cellular retinoic acid-binding protein
CRBP	cellular retinol-binding protein
CV	coefficient of variation
CYP	cytochrome P450
D-A-CH	Deutschland-Austria-Confoederatio Helvetica
DGAT	acyl-CoA:retinol acyltransferase
DH	UK Department of Health
DIPP	type 1 Diabetes Prediction and Prevention survey
DNFCS	Dutch National Food Consumption Survey
DRD	deuterated-retinol-dilution
DRV	Dietary Reference Value
EAR	Estimated Average Requirement
EFSA	European Food Safety Authority
EsKiMo	Ernährungstudie als KIGGS-Modul
EU	European Union
FABP	fatty acid-binding protein
FC_PREGNANTWOMEN	food consumption of pregnant women in Latvia
FAO	Food and Agriculture Organization of the United Nations
FINDIET	the national dietary survey of Finland
FFQ	food frequency questionnaire

HDL	high-density lipoprotein
HR	hazard ratio
IFN	interferon
IL	interleukin
INCA	étude Individuelle Nationale de Consommations Alimentaires
INRAN-SCAI	Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione - Studio sui Consumi Alimentari in Italia
IOM	US Institute of Medicine of the National Academy of Sciences
LDL	low-density lipoprotein
LRAT	lecithin:retinol acyltransferase
NANS	National Adult Nutrition Survey
NDNS	UK National Diet and Nutrition Survey
NHANES III	US Third National Health and Nutrition Examination Survey
NNR	Nordic Nutrition Recommendations
NPC	Nutritional Prevention of Cancer
NWSSP	Nutrition and Wellbeing of Secondary School Pupils
PPAR	peroxisome proliferator-activated receptor
PRI	Population Reference Intake
RAE	retinol activity equivalent
RAR	retinoic acid receptor
RBP	retinol-binding protein
RDA	Recommended Dietary Allowance
RDR	relative dose response
RE	retinol equivalent
RID	retinol isotope dilution
RNI	Reference Nutrient Intake
RXR	retinoic X receptor
SACN	UK Scientific Advisory Committee on Nutrition

SCF	Scientific Committee for Food
SD	standard deviation
SE	standard error
SEM	standard error of the mean
SR-B	scavenger receptor class B
UK	United Kingdom
UL	Tolerable Upper Intake Level
UNU	United Nations University
US	United States
VELS	Verzehrsstudie zur Ermittlung der Lebensmittelaufnahme von Säuglingen und Kleinkindern für die Abschätzung eines akuten Toxizitätsrisikos durch Rückstände von Pflanzenschutzmitteln
VLDL	very low-density lipoprotein
WHAS	Women's Health and Ageing Study
WHO	World Health Organization